

Site report 2018 Kutupalong New BANGLADESH



Population


25,071 refugees

15%

Children under 5

20%

Women of reproductive age



Origin of refugees

MYANMAR

Site opened
Aug 2017

Partners



Save the Children



Key Indicators

CRUDE MORTALITY RATE

0.2

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.5

Deaths/1,000 refugees/month

MATERNAL MORTALITY

2

Deaths

SKILLED BIRTH ATTENDANCE

83%

of all births

GAM

NC

Global Acute Malnutrition

SAM

NC

Severe Acute Malnutrition

WATER SUPPLY

17


Liters/ person/ day

LATRINE COVERAGE



2

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Number of medical doctors	52	1 : < 50,000	✓	Incidence of malaria among children under 5	0	
Number of qualified nurses	67	1 : < 10,000	✓	Incidence of watery diarrhoea among children under 5	22	
Number of community health workers	302	1 : < 1,000	✓	Incidence of pneumonia among children under 5	20	
ACCESS AND UTILIZATION			VACCINATION			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Consultations per trained clinician per day	27	< 50	✓	Incidence of pneumonia among over 5	6	
Health utilisation rate (new visits / refugee / year)	0.9	1 - 4	⚠	Tuberculosis success rate	89% < 90% ⚠	
Proportion of host population consultations	0%			Were any MDR/X-TB cases diagnosed among PoCs?	No No ✗	
MORTALITY			IN-PATIENT DEPARTMENT (IPD)			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75	✓	Full vaccination coverage	0% < 95% ✗	
Under-five Mortality Rate (U5MR) (/1000/month)	0.5	< 1.5	✓	Measles vaccination coverage	Not collected	
Infant Mortality Rate (IMR) (/1000 livebirths)	48.9	< 30	✗	Source of vaccination data	survey	
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	22	< 20	⚠	IN-PATIENT DEPARTMENT (IPD)		
OUTBREAK ALERT AND RESPONSE				INDICATOR	STANDARD	
	INDICATOR	STANDARD				
Proportion of outbreaks investigated within 48 hours		100%		Average length of stay (days)	5.3	
Number of outbreaks reported	0			Case fatality rate	0.0	
				Hospitalisation rate	14.6 50 - 150 ✗	

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Antenatal care coverage	94%	> 90%	✓	Contraceptive prevalence rate	78% ≥ 30% ✓	
Coverage of antenatal tetanus vaccination	75%	> 95%	⚠	Source of contraceptive prevalence data	HIS	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	83%	≥ 90%	⚠	Total number of reported rapes	0	
Proportion of births conducted by caesarean section	0%	5 - 15%	✗	Proportion of eligible rape survivors provided with PEP within 72 hours	100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	2%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%	
Proportion of teenage pregnancies	2%			SEXUALLY TRANSMITTED INFECTIONS (STIS)		
Still birth rate (/ 1000 total births / month)	2				INDICATOR	STANDARD
Number of maternal deaths	2			Number of genital ulcer diseases	6	
Proportion of maternal deaths investigated within 48 hours	0%	100%	✗	Number of Pelvic Inflammatory diseases	0	

HIV/AIDS

PREVENTION & TREATMENT

Proportion of blood units screened for HIV

INDICATOR STANDARD

100%

Condom distribution rate

0.02 > 0.5

Proportion of TB patients tested for HIV

100%

PREVENTION & TREATMENT

Number of PoCs receiving ART

0

✘ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

0% 100%



Nutrition

NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

0% > 70%



CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

INDICATOR STANDARD

Recovery rate

> 75%

Default rate

< 0%

Death rate

< 10%

Referral rate

MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

INDICATOR STANDARD

Recovery rate

> 75%

Default rate

< 15%

Death rate

< 3%

Referral rate

Water, Sanitation and Hygiene (WASH)

WATER

Average number of litres of potable water available per person per day

INDICATOR STANDARD

17 < 20

Proportion of households collecting >= 15 litres per person per day

90% > 95%

Number of persons of concern per water tap

67 < 80

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

85% > 95%

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml

94% > 95%

Proportion of households collecting drinking water from protected water sources only

95% > 95%

Proportion of households with sufficient daily water storage capacity

57% > 80%

WATER STAFFING

Number of hygiene promoters

INDICATOR STANDARD

233 1 : < 20,000

SANITATION

Refugees per latrine/toilet

INDICATOR STANDARD

2 < 20

Proportion of households with drop-hole latrine or drop-hole toilet

Proportion of households report defecating in a toilet

91% > 85%

Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

83% > 90%