

Site report 2018 Nayapara BANGLADESH

Population

19,895 refugees







Origin of refugees MYANMAR

Site opened Jan 1992

Partners



















Key Indicators

CRUDE MORTALITY RATE

GAM

Deaths/1,000 refugees/ month

month

SAM

Global Acute 1%

U5 MORTALITY RATE

Deaths/1,000 refugees/

Severe Acute

MATERNAL MORTALITY

Deaths

WATER SUPPLY

Liters/ person/ day


SKILLED BIRTH ATTENDANCE


100% of all births


LATRINE COVERAGE

Refugees/communal


COMMUNICABLE DISEASES








NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD		
Number of medical doctors	4	1:<50,000	•	Incidence of malaria among children under 5	0			
Number of qualified nurses	8	1:<10,000	②	Incidence of watery diarrhoea among	34			
Number of community health workers	26	1 : < 1,000	②	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	164			
Consultations per trained clinician per day	40	< 50	②	Incidence of pneumonia among over 5	34			
Health utilisation rate (new visits / refugee /	2.9	1 - 4	•	Tuberculosis success rate	87%	< 90%	A	
year) Proportion of host population consultations	0%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	©	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	STAN	DARD	
Crude Mortality Rate (CMR) (/1000/month)	0.4	< 0.75	②	Full vaccination coverage	78%	< 9	5%	8
Under-five Mortality Rate (U5MR) (/1000/month)	0.8	< 1.5	②	Measles vaccination coverage		Not collected	d	
Infant Mortality Rate (IMR) (/1000 livebirths)	30.5	< 30	A	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	20	< 20	<u> </u>	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STAN	DARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	3.9			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	73.4	50 -	150	②

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	79%	> 90%	8	Contraceptive prevalence rate	73%	≥ 30%	•
Coverage of antenatal tetanus vaccination	79%	> 95%	<u> </u>	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	100%	≥ 90%	②	Total number of reported rapes	6		
Proportion of births conducted by caesarean section	2%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours	20%	100%	8
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	5%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours	80%	100%	×
Proportion of teenage pregnancies	9%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	2			Number of genital ulcer diseases	9		
Number of maternal deaths	2			Number of Pelvic Inflammatory diseases	0		
Proportion of maternal deaths investigated within 48 hours	50%	100%	8				

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receving ART	0	
Condom distribution rate	0.08	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	0%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	12%	< 20%	A	Proportion of children 0-23 months who were put to the breast within one hour of birth	Not	collected	
Prevalence of stunting (6-59 months)	38%	< 20%	②	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not	collected	
Prevalence of severe acute malnutrition (6 - 59 months)	1%	< 3%	②	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	38%	< 20%	②	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not	collected		Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	95%	> 70%	Ø
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	86			Number of new moderate acute admissions (6-59 months)	613		
Recovery rate	94%	> 75%	②	Recovery rate	67%	> 75%	A
Default rate	0%	< 0%	8	Default rate	2%	< 15%	②
Death rate	0%	< 10%	②	Death rate	0%	< 3%	②
Referral rate	1%			Referral rate	14%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	11	< 20	8	Number of hygiene promoters	106	1:<20,000	•
Proportion of households collecting >= 15 litres per person per day	0%	> 95%	8	SANITATION Refugees per latrine/toilet	INDICATOR 5	STANDARD < 20	©
Number of persons of concern per water tap	106	< 80	8	Proportion of households with drop-hole	3	< 20	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%	•	latrine or drop-hole toilet Proportion of households report defecating in a toilet	92%	> 85%	•
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	100%	> 95%	•	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	86%	> 90%	A
Proportion of households collecting drinking water from protected water sources only	87%	> 95%	A				
Proportion of households with sufficient daily water storage capacity	83%	> 80%	•				