



Site report 2018 Goudebo BURKINA FASO



Population

7,768 refugees



Origin of refugees

MALI CENTRAL AFRICAN
REPUBLIC CHAD BURUNDI

Site opened
Feb 2012

Partners

CSSI



Key Indicators

CRUDE MORTALITY RATE

0.3 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.7 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

100% of all births

GAM

NC Global Acute Malnutrition

SAM

NC Severe Acute Malnutrition


WATER SUPPLY

15 Liters/ person/ day


LATRINE COVERAGE

8 Refugees/communal toilets or latrines



COMMUNICABLE DISEASES




NON-COMMUNICABLE DISEASES



INJURIES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	1	1 : < 50,000	✓	Incidence of malaria among children under 5	39
Number of qualified nurses	7	1 : < 10,000	✓	Incidence of watery diarrhoea among children under 5	0
Number of community health workers	18	1 : < 1,000	✓	Incidence of pneumonia among children under 5	57
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	10	< 50	✓	Incidence of pneumonia among over 5	28
Health utilisation rate (new visits / refugee / year)	2.0	1 - 4	✓	Tuberculosis success rate	100% < 90% ✓
Proportion of host population consultations	12%			Were any MDR/X-TB cases diagnosed among PoCs?	No No ✗
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.3	< 0.75	✓	Full vaccination coverage	98% < 95% ✓
Under-five Mortality Rate (U5MR) (/1000/month)	0.7	< 1.5	✓	Measles vaccination coverage	107% < 95% ✓
Infant Mortality Rate (IMR) (/1000 livebirths)	29.8	< 30	✓	Source of vaccination data	HIS
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	✓	Average length of stay (days)	2.5
OUTBREAK ALERT AND RESPONSE			CASE FATALITY RATE		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours	100%	100%	✓	Case fatality rate	0.0
Number of outbreaks reported	2			Hospitalisation rate	61.5 50 - 150 ✓

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	37%	> 90%	✗	Contraceptive prevalence rate	36% ≥ 30% ✓
Coverage of antenatal tetanus vaccination	78%	> 95%	⚠	Source of contraceptive prevalence data	HIS
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	100%	≥ 90%	✓	Total number of reported rapes	0
Proportion of births conducted by caesarean section	1%	5 - 15%		Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	5%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%
Proportion of teenage pregnancies	5%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	16
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	20
Proportion of maternal deaths investigated within 48 hours		100%			

HIV/AIDS

PREVENTION & TREATMENT

Proportion of blood units screened for HIV

Condom distribution rate

Proportion of TB patients tested for HIV

INDICATOR

STANDARD

100%

0.03 > 0.5

100%

PREVENTION & TREATMENT

Number of PoCs receiving ART

✘ Proportion of HIV positive mothers who swallowed ARV during delivery

PMTCT coverage

INDICATOR

STANDARD

9

100%

100%

100%



Nutrition

NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

Prevalence of stunting (6-59 months)

Prevalence of severe acute malnutrition (6 - 59 months)

Prevalence of anaemia in children (6-59 months)

Prevalence of anaemia in women of reproductive age (15-49 yrs)

INDICATOR

STANDARD

Not collected

Not collected

Not collected

Not collected

Not collected

INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

IYCF programmes targeting pregnant and lactating women established or maintained?

INDICATOR

STANDARD

Not collected

Not collected

Yes

MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

INDICATOR

STANDARD

Not collected

CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

Recovery rate

Default rate

Death rate

Referral rate

INDICATOR

STANDARD

> 75%

< 0%

< 10%

MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

Recovery rate

Default rate

Death rate

Referral rate

INDICATOR

STANDARD

77

100%

0%

0%

0%

> 75%

< 15%

< 3%



Water, Sanitation and Hygiene (WASH)

WATER

Average number of litres of potable water available per person per day

Proportion of households collecting >= 15 litres per person per day

Number of persons of concern per water tap

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml

Proportion of households collecting drinking water from protected water sources only

Proportion of households with sufficient daily water storage capacity

INDICATOR

STANDARD

15

100%

100

100%

100%

100%

100%

< 20

> 95%

< 80

> 95%

> 95%

> 95%

> 80%

WATER STAFFING

✘ Number of hygiene promoters

SANITATION

Refugees per latrine/toilet

⚠ Proportion of households with drop-hole latrine or drop-hole toilet

✔ Proportion of households report defecating in a toilet

✔ Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

INDICATOR

STANDARD

4

8

100%

85%

77%

1 : < 20,000

< 20

> 85%

> 90%



WATER

Average number of litres of potable water available per person per day

Proportion of households collecting >= 15 litres per person per day

Number of persons of concern per water tap

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml

Proportion of households collecting drinking water from protected water sources only

Proportion of households with sufficient daily water storage capacity

INDICATOR

STANDARD

15

100%

100

100%

100%

100%

100%

< 20

> 95%

< 80

> 95%

> 95%

> 95%

> 80%

WATER STAFFING

✘ Number of hygiene promoters

SANITATION

Refugees per latrine/toilet

⚠ Proportion of households with drop-hole latrine or drop-hole toilet

✔ Proportion of households report defecating in a toilet

✔ Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

INDICATOR

STANDARD

4

8

100%

85%

77%

1 : < 20,000

< 20

> 85%

> 90%

