

Site report 2018 Mentao BURKINA FASO



Population


6,624 refugees

17%

Children under 5

20%

Women of reproductive age



Origin of refugees

MALI

Site opened

Feb 2012

Partners

CSSI



Key Indicators

CRUDE MORTALITY RATE

0.2

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.5

Deaths/1,000 refugees/month

MATERNAL MORTALITY

0

Deaths

SKILLED BIRTH ATTENDANCE

96%

of all births

GAM

NC

Global Acute Malnutrition

SAM

NC

Severe Acute Malnutrition

WATER SUPPLY

22


Liters/ person/ day

LATRINE COVERAGE



4

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	1	1 : < 50,000	✓	Incidence of malaria among children under 5	56
Number of qualified nurses	7	1 : < 10,000	✓	Incidence of watery diarrhoea among children under 5	0
Number of community health workers	19	1 : < 1,000	✓	Incidence of pneumonia among children under 5	109
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	12	< 50	✓	Incidence of pneumonia among over 5	47
Health utilisation rate (new visits / refugee / year)	1.7	1 - 4	✓	Tuberculosis success rate	100% < 90% ✓
Proportion of host population consultations	51%			Were any MDR/X-TB cases diagnosed among PoCs?	No No ✗
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75	✓	Full vaccination coverage	92% < 95% ⚠
Under-five Mortality Rate (U5MR) (/1000/month)	0.5	< 1.5	✓	Measles vaccination coverage	89% < 95% ⚠
Infant Mortality Rate (IMR) (/1000 livebirths)	43.9	< 30	✗	Source of vaccination data	HIS
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	18	< 20	✓	Average length of stay (days)	2.2
OUTBREAK ALERT AND RESPONSE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0
Number of outbreaks reported	0			Hospitalisation rate	56.3 50 - 150 ✓

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	43%	> 90%	✗	Contraceptive prevalence rate	53% ≥ 30% ✓
Coverage of antenatal tetanus vaccination	88%	> 95%	⚠	Source of contraceptive prevalence data	HIS
DELIVERY CARE			SEXUALLY TRANSMITTED INFECTIONS (STIS)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	96%	≥ 90%	✓	Total number of reported rapes	0
Proportion of births conducted by caesarean section	1%	5 - 15%	✗	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	11%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%
Proportion of teenage pregnancies	6%			Number of genital ulcer diseases	3
Still birth rate (/ 1000 total births / month)	0			Number of Pelvic Inflammatory diseases	4
Number of maternal deaths	0			Proportion of maternal deaths investigated within 48 hours	100%

HIV/AIDS

PREVENTION & TREATMENT

Proportion of blood units screened for HIV

INDICATOR STANDARD

100%

Condom distribution rate

0.01

> 0.5

Proportion of TB patients tested for HIV

100%

PREVENTION & TREATMENT

Number of PoCs receiving ART

9

✘ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

101%

100%



Nutrition

NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

INDICATOR STANDARD

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

Not collected

CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

INDICATOR STANDARD

Recovery rate

> 75%

Default rate

< 0%

Death rate

< 10%

Referral rate

MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

92

Recovery rate

100%

> 75%



Default rate

0%

< 15%



Death rate

0%

< 3%



Referral rate

0%

Water, Sanitation and Hygiene (WASH)

WATER

Average number of litres of potable water available per person per day

INDICATOR STANDARD

22

< 20



Proportion of households collecting >= 15 litres per person per day

100%

> 95%



Number of persons of concern per water tap

70

< 80



Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

100%

> 95%



Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/100 ml

100%

> 95%



Proportion of households collecting drinking water from protected water sources only

100%

> 95%



Proportion of households with sufficient daily water storage capacity

100%

> 80%



WATER STAFFING

Number of hygiene promoters

INDICATOR STANDARD

6

1 : < 20,000



SANITATION

Refugees per latrine/toilet

INDICATOR STANDARD

4

< 20



Proportion of households with drop-hole latrine or drop-hole toilet

100%

Proportion of households report defecating in a toilet

85%

> 85%



Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

77%

> 90%

