

Site report 2018 Musasa BURUNDI



Population


8,420 refugees

19%

Children under 5

20%

Women of reproductive age



Origin of refugees

DRC **RWA** **ETIO** **SOM**

Site opened
Jan 2007

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.3 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

100% of all births

GAM

NC Global Acute Malnutrition

SAM

NC Severe Acute Malnutrition


WATER SUPPLY

24 Liters/ person/ day



LATRINE COVERAGE

22 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000	⊖	Incidence of malaria among children under 5	27
Number of qualified nurses		1 : < 10,000	⊖	Incidence of watery diarrhoea among children under 5	23
Number of community health workers		1 : < 1,000	⊖	Incidence of pneumonia among children under 5	101
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	42	< 50	✓	Incidence of pneumonia among over 5	43
Health utilisation rate (new visits / refugee / year)	3.1	1 - 4	✓	Tuberculosis success rate	< 90%
Proportion of host population consultations	2%			Were any MDR/X-TB cases diagnosed among PoCs?	No
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	✓	Full vaccination coverage	43% < 95% ✗
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5	✓	Measles vaccination coverage	50% < 95% ✗
Infant Mortality Rate (IMR) (/1000 livebirths)	15.5	< 30	✓	Source of vaccination data	
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	✓	Average length of stay (days)	3.9
OUTBREAK ALERT AND RESPONSE			CASE FATALITY RATE		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0
Number of outbreaks reported	0			Hospitalisation rate	82.1 50 - 150 ✓

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	100%	> 90%	✓	Contraceptive prevalence rate	24% ≥ 30% ✗
Coverage of antenatal tetanus vaccination	80%	> 95%	⚠	Source of contraceptive prevalence data	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	100%	≥ 90%	✓	Total number of reported rapes	1
Proportion of births conducted by caesarean section	7%	5 - 15%	✓	Proportion of eligible rape survivors provided with PEP within 72 hours	100% 100% ✓
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	7%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	0% 100% ✗
Proportion of teenage pregnancies	0%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	41
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0
Proportion of maternal deaths investigated within 48 hours		100%			

HIV/AIDS

PREVENTION & TREATMENT

Proportion of blood units screened for HIV

INDICATOR STANDARD

100%

Condom distribution rate

0.83 > 0.5

Proportion of TB patients tested for HIV

100%

PREVENTION & TREATMENT

Number of PoCs receiving ART

INDICATOR STANDARD

14

✓ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

94% 100%



Nutrition

NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

INDICATOR STANDARD

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

INDICATOR STANDARD

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

INDICATOR STANDARD

> 70%

CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

INDICATOR STANDARD

Recovery rate

> 75%

Default rate

< 0%

Death rate

< 10%

Referral rate

MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

INDICATOR STANDARD

55

Recovery rate

97% > 75%



Default rate

0% < 15%



Death rate

0% < 3%



Referral rate

3%

Water, Sanitation and Hygiene (WASH)

WATER

Average number of litres of potable water available per person per day

INDICATOR STANDARD

24 < 20

Proportion of households collecting >= 15 litres per person per day

> 95%

Number of persons of concern per water tap

< 80

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

> 95%

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml

> 95%

Proportion of households collecting drinking water from protected water sources only

> 95%

Proportion of households with sufficient daily water storage capacity

> 80%

WATER STAFFING

Number of hygiene promoters

INDICATOR STANDARD

1 : < 20,000

SANITATION

Refugees per latrine/toilet

INDICATOR STANDARD

22 < 20



Proportion of households with drop-hole latrine or drop-hole toilet

Proportion of households report defecating in a toilet

> 85%

Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

> 90%