

Site report 2018 Bwagiriza BURUNDI



Population


9,646 refugees

16%

Children under 5

20%

Women of reproductive age



Origin of refugees

DRC **RWA** **SOMALIA** **SOMALIE**

Site opened
May 2009

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.2 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

100% of all births

GAM

NC Global Acute Malnutrition

SAM

NC Severe Acute Malnutrition


WATER SUPPLY

23 Liters/ person/ day



LATRINE COVERAGE

5 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000
Number of qualified nurses		1 : < 10,000
Number of community health workers		1 : < 1,000

ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	54	< 50
Health utilisation rate (new visits / refugee / year)	3.3	1 - 4
Proportion of host population consultations	6%	

MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20

OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%
Number of outbreaks reported	0	

MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	37	
Incidence of watery diarrhoea among children under 5	54	
Incidence of pneumonia among children under 5	45	
Incidence of pneumonia among over 5	22	
Tuberculosis success rate		< 90%
Were any MDR/X-TB cases diagnosed among PoCs?		No

VACCINATION

	INDICATOR	STANDARD	
Full vaccination coverage	0%	< 95%	✗
Measles vaccination coverage	116%	< 95%	✓
Source of vaccination data			✓

IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD	
Average length of stay (days)	3.4		
Case fatality rate	0.0		
Hospitalisation rate	67.7	50 - 150	✓

Reproductive Health

ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	96%	> 90%
Coverage of antenatal tetanus vaccination	88%	> 95%

DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	100%	≥ 90%
Proportion of births conducted by caesarean section	7%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	3%	< 15%
Proportion of teenage pregnancies	9%	
Still birth rate (/ 1000 total births / month)	0	
Number of maternal deaths	0	
Proportion of maternal deaths investigated within 48 hours		100%

FAMILY PLANNING

	INDICATOR	STANDARD	
Contraceptive prevalence rate	11%	≥ 30%	✗
Source of contraceptive prevalence data			⚠

SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD	
Total number of reported rapes	0		✓
Proportion of eligible rape survivors provided with PEP within 72 hours		100%	✓
Proportion of eligible rape survivors provided with ECP within 120 hours		100%	✓

SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD	
Number of genital ulcer diseases	29		
Number of Pelvic Inflammatory diseases	0		

HIV/AIDS

PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.46	> 0.5
Proportion of TB patients tested for HIV		100%

PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	40		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	⚠️
PMTCT coverage	92%	100%	⚠️

Nutrition

NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	Not collected	
Prevalence of stunting (6-59 months)	Not collected	
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected	
Prevalence of anaemia in children (6-59 months)	Not collected	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected	

INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD
Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected	
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected	
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes

MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%

CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)		
Recovery rate		> 75%
Default rate		< 0%
Death rate		< 10%
Referral rate		

MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	81		
Recovery rate	74%	> 75%	⚠️
Default rate	5%	< 15%	✅
Death rate	0%	< 3%	✅
Referral rate	1%		

Water, Sanitation and Hygiene (WASH)

WATER

	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	23	< 20	⚠️
Proportion of households collecting >= 15 litres per person per day		> 95%	
Number of persons of concern per water tap		< 80	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	
Proportion of households collecting drinking water from protected water sources only		> 95%	
Proportion of households with sufficient daily water storage capacity		> 80%	

WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters		1 : < 20,000	⊖

SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	5	< 20	✅
Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of households report defecating in a toilet		> 85%	
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	