

# Site report 2018 Kavumu BURUNDI

### **Population**

18,964 refugees







#### **Origin of refugees**

TANZANIA

**SOMALIA** 

Site opened **Apr 2015** 

#### **Partners**





### **Key Indicators**

**CRUDE MORTALITY RATE** 

Deaths/1,000 refugees/

**GAM** 

NC Global Acute U5 MORTALITY RATE

Deaths/1,000 refugees/

SAM

NC

Severe Acute Malnutrition

MATERNAL MORTALITY

Deaths

WATER SUPPLY

18

Liters/ person/ day

SKILLED BIRTH **ATTENDANCE** 

92%


of all births


LATRINE COVERAGE

10


Refugees/communal toilets or latrines

#### COMMUNICABLE DISEASES





### **NON-COMMUNICABLE DISEASES**



## Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STA	NDARD	
Number of medical doctors		1:<50,000	•	Incidence of malaria among children under 5	33			
Number of qualified nurses		1 : < 10,000	•	Incidence of watery diarrhoea among	29			
Number of community health workers		1:<1,000	•	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	13			
Consultations per trained clinician per day	85	< 50	8	Incidence of pneumonia among over 5	7			
Health utilisation rate (new visits / refugee /	2.4	1 - 4	•	Tuberculosis success rate		<	90%	
year)	40/			Were any MDR/X-TB cases diagnosed among PoCs?			No	
Proportion of host population consultations	4%							
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	OR	STANDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75	<b>②</b>	Full vaccination coverage	8%		< 95%	•
Under-five Mortality Rate (U5MR) (/1000/month)	0.6	< 1.5	<b>②</b>	Measles vaccination coverage	14%		< 95%	•
Infant Mortality Rate (IMR) (/1000 livebirths)	42.0	< 30	8	Source of vaccination data				
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	14	< 20	<b>⊘</b>	IN-PATIENT DEPARTMENT (IPD)	INDICATO	OR	STANDARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	2.7			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	36.8		50 - 150	•

# Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	59%	> 90%	8	Contraceptive prevalence rate	4%	≥ 30%	E
Coverage of antenatal tetanus vaccination	18%	> 95%	8	Source of contraceptive prevalence data			
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	92%	≥ 90%	<b>②</b>	Total number of reported rapes	6		
Proportion of births conducted by caesarean section	4%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours	0%	100%	3
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	4%	< 15%	<b>②</b>	Proportion of eligible rape survivors provided with ECP within 120 hours	83%	100%	3
Proportion of teenage pregnancies	1%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate ( / 1000 total births / month)	0			Number of genital ulcer diseases	228		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	13		
Proportion of maternal deaths investigated within 48 hours		100%					

### **HIV/AIDS**

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	93	
Condom distribution rate	0.81	> 0.5	<b>②</b>	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	98%	100%

## Nutrition

NUTRITION GENERAL	INDICATOR STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected	Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	Not collected	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%	
CMAM: SAM TREATMENT	INDICATOR STANDARD	MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)		Number of new moderate acute admissions (6-59 months)	239		
Recovery rate	> 75%	Recovery rate	77%	> 75%	<b>Ø</b>
Default rate	< 0%	Default rate	8%	< 15%	<b>Ø</b>
Death rate	< 10%	Death rate	0%	< 3%	<b>Ø</b>
Referral rate		Referral rate	8%		

# Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	18	< 20	A	Number of hygiene promoters		1:<20,000	•
Proportion of households collecting >= 15 litres per person per day		> 95%		SANITATION  Refugees per latrine/toilet	INDICATOR 10	STANDARD < 20	<b>②</b>
Number of persons of concern per water tap		< 80		Proportion of households with drop-hole		- 20	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5		> 95%		Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%		Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	
Proportion of households collecting drinking water from protected water sources only		> 95%					
Proportion of households with sufficient daily water storage capacity		> 80%					