

Site report 2018 Bili CONGO (DR)



Population

9,381 refugees

24%

Children under 5

20%

Women of reproductive age



Origin of refugees

CENTRAL AFRICAN REPUBLIC

Site opened

Mar 2015

Partners



Key Indicators

CRUDE MORTALITY RATE

0.2

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.4

Deaths/1,000 refugees/month

MATERNAL MORTALITY

0

Deaths

SKILLED BIRTH ATTENDANCE

99%

of all births

GAM

NC

Global Acute Malnutrition

SAM

NC

Severe Acute Malnutrition

WATER SUPPLY

19


Liters/ person/ day

LATRINE COVERAGE



22

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	1	1 : < 50,000	✓	139	
Number of qualified nurses	10	1 : < 10,000	✓	6	
Number of community health workers	12	1 : < 1,000	✓	47	
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	21	< 50	✓	67%	< 95%
Health utilisation rate (new visits / refugee / year)	1.8	1 - 4	✓	68%	< 95%
Proportion of host population consultations	10%		✓	HIS	
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75	✓	Average length of stay (days)	3.0
Under-five Mortality Rate (U5MR) (/1000/month)	0.4	< 1.5	✓	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	10.4	< 30	✓	Hospitalisation rate	91.8
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	✓		50 - 150
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%			
Number of outbreaks reported	0				

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	89%	> 90%	⚠	28%	≥ 30%
Coverage of antenatal tetanus vaccination	92%	> 95%	⚠	Source of contraceptive prevalence data	HIS
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	99%	≥ 90%	✓	Total number of reported rapes	7
Proportion of births conducted by caesarean section	3%	5 - 15%	✗	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	6%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%
Proportion of teenage pregnancies	2%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	64
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	59
Proportion of maternal deaths investigated within 48 hours		100%			

HIV/AIDS

PREVENTION & TREATMENT

Proportion of blood units screened for HIV

INDICATOR STANDARD

100%

Condom distribution rate

0.62 > 0.5

Proportion of TB patients tested for HIV

100%

PREVENTION & TREATMENT

Number of PoCs receiving ART

INDICATOR STANDARD

26

✓ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

80% 100%



Nutrition

NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

INDICATOR STANDARD

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

INDICATOR STANDARD

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

INDICATOR STANDARD

Not collected

CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

INDICATOR STANDARD

48

Recovery rate

100% > 75%

Default rate

0% < 0%

Death rate

0% < 10%

Referral rate

0%

MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

INDICATOR STANDARD

68

✓ Recovery rate

69% > 75%



✗ Default rate

9% < 15%



✓ Death rate

0% < 3%



Referral rate

22%

Water, Sanitation and Hygiene (WASH)

WATER

Average number of litres of potable water available per person per day

INDICATOR STANDARD

19 < 20

Proportion of households collecting >= 15 litres per person per day

95% > 95%

Number of persons of concern per water tap

150 < 80

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

90% > 95%

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/100 ml

0% > 95%

Proportion of households collecting drinking water from protected water sources only

100% > 95%

Proportion of households with sufficient daily water storage capacity

50% > 80%

WATER STAFFING

Number of hygiene promoters

INDICATOR STANDARD

7 1 : < 20,000

SANITATION

Refugees per latrine/toilet

INDICATOR STANDARD

22 < 20

✗ Proportion of households with drop-hole latrine or drop-hole toilet

65%

Proportion of households report defecating in a toilet

69% > 85%

✗ Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

90% > 90%