

Site report 2018 Boyabo CONGO (DR)



Population


18,344 refugees

21%

Children under 5

20%

Women of reproductive age



Origin of refugees

CENTRAL AFRICAN REPUBLIC

Site opened
Jun 2013

Partners



Key Indicators

CRUDE MORTALITY RATE

0.2

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.3

Deaths/1,000 refugees/month

MATERNAL MORTALITY

0

Deaths

SKILLED BIRTH ATTENDANCE

100%

of all births

GAM

NC

Global Acute Malnutrition

SAM

NC

Severe Acute Malnutrition

WATER SUPPLY

20


Liters/ person/ day

LATRINE COVERAGE



16

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors	1	1 : < 50,000
Number of qualified nurses	12	1 : < 10,000
Number of community health workers	15	1 : < 1,000

ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	18	< 50
Health utilisation rate (new visits / refugee / year)	0.9	1 - 4
Proportion of host population consultations	7%	

MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	7.4	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20

OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours	0%	100%
Number of outbreaks reported	1	

MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	92	
Incidence of watery diarrhoea among children under 5	4	
Incidence of pneumonia among children under 5	9	
Incidence of pneumonia among over 5	3	
Tuberculosis success rate	63%	< 90%
Were any MDR/X-TB cases diagnosed among PoCs?	No	No

VACCINATION

	INDICATOR	STANDARD
Full vaccination coverage	73%	< 95%
Measles vaccination coverage	73%	< 95%
Source of vaccination data	HIS	

IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD
Average length of stay (days)	1.8	
Case fatality rate	0.0	
Hospitalisation rate	24.6	50 - 150

Reproductive Health

ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	75%	> 90%
Coverage of antenatal tetanus vaccination	89%	> 95%

DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	100%	≥ 90%
Proportion of births conducted by caesarean section	2%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	5%	< 15%
Proportion of teenage pregnancies	3%	
Still birth rate (/ 1000 total births / month)	3	
Number of maternal deaths	0	
Proportion of maternal deaths investigated within 48 hours		100%

FAMILY PLANNING

	INDICATOR	STANDARD
Contraceptive prevalence rate	5%	≥ 30%
Source of contraceptive prevalence data	HIS	

SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD
Total number of reported rapes	17	
Proportion of eligible rape survivors provided with PEP within 72 hours	100%	100%
Proportion of eligible rape survivors provided with ECP within 120 hours	100%	100%

SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD
Number of genital ulcer diseases	36	
Number of Pelvic Inflammatory diseases	40	

HIV/AIDS

PREVENTION & TREATMENT

Proportion of blood units screened for HIV

100%

100%

Condom distribution rate

0.58

> 0.5

Proportion of TB patients tested for HIV

100%

PREVENTION & TREATMENT

✓ Number of PoCs receiving ART

-41

✓ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

92%

100%



Nutrition

NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

Not collected

CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

17

Recovery rate

50%

> 75%

Default rate

0%

< 0%

Death rate

50%

< 10%

Referral rate

0%

MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

90

✗ Recovery rate

> 75%

✗ Default rate

< 15%

✗ Death rate

< 3%

Referral rate

Water, Sanitation and Hygiene (WASH)

WATER

Average number of litres of potable water available per person per day

20

< 20



Proportion of households collecting >= 15 litres per person per day

Not collected

Number of persons of concern per water tap

Not collected

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

Not collected

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml

Not collected

Proportion of households collecting drinking water from protected water sources only

Not collected

Proportion of households with sufficient daily water storage capacity

Not collected

WATER STAFFING

Number of hygiene promoters

5

1 : < 20,000



SANITATION

Refugees per latrine/toilet

16

< 20



Proportion of households with drop-hole latrine or drop-hole toilet

Not collected

Proportion of households report defecating in a toilet

Not collected

Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

Not collected

INDICATOR

STANDARD

INDICATOR

STANDARD