

# Site report 2018 Lusenda CONGO (DR)



## Population

**28,202** refugees

**22%**

Children under 5

**20%**

Women of reproductive age



Origin of refugees

**BURUNDI**

Site opened

**Apr 2015**

## Partners



## Key Indicators

CRUDE MORTALITY RATE

**0.1** Deaths/1,000 refugees/month

U5 MORTALITY RATE

**0.3** Deaths/1,000 refugees/month

MATERNAL MORTALITY

**3** Deaths

SKILLED BIRTH ATTENDANCE

**100%** of all births

GAM

**NC** Global Acute Malnutrition

SAM

**NC** Severe Acute Malnutrition


WATER SUPPLY

**21** Liters/ person/ day



LATRINE COVERAGE

**15** Refugees/communal toilets or latrines


### COMMUNICABLE DISEASES




### INJURIES



### NON-COMMUNICABLE DISEASES



### MENTAL HEALTH



## Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	2	1 : < 50,000	✓	72	
Number of qualified nurses	25	1 : < 10,000	✓	13	
Number of community health workers	25	1 : < 1,000	⚠	10	
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	41	< 50	✓	29%	< 95%
Health utilisation rate (new visits / refugee / year)	1.4	1 - 4	✓	34%	< 95%
Proportion of host population consultations	0%		✓	HIS	
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	✓	Average length of stay (days)	0.3
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5	✓	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	15.8	< 30	✓	Hospitalisation rate	282.4
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	5	< 20	✓		50 - 150
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD			
Proportion of outbreaks investigated within 48 hours		100%			
Number of outbreaks reported	0				

## Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	79%	> 90%	✗	53%	≥ 30%
Coverage of antenatal tetanus vaccination	63%	> 95%	⚠	HIS	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	100%	≥ 90%	✓	Total number of reported rapes	25
Proportion of births conducted by caesarean section	8%	5 - 15%	✓	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	5%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	7%
Proportion of teenage pregnancies	0%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate ( / 1000 total births / month)	2			Number of genital ulcer diseases	101
Number of maternal deaths	3			Number of Pelvic Inflammatory diseases	10
Proportion of maternal deaths investigated within 48 hours	100%	100%	✓		

## HIV/AIDS

### PREVENTION & TREATMENT

Proportion of blood units screened for HIV

INDICATOR STANDARD

100%

Condom distribution rate

0.25

> 0.5

Proportion of TB patients tested for HIV

100%

### PREVENTION & TREATMENT

Number of PoCs receiving ART

INDICATOR STANDARD

264

✘ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

97%

100%



## Nutrition

### NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

INDICATOR STANDARD

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

### INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

INDICATOR STANDARD

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

### MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

INDICATOR STANDARD

Not collected

### CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

INDICATOR STANDARD

352

Recovery rate

98%

> 75%



Default rate

0%

< 0%



Death rate

0%

< 10%



Referral rate

2%

### MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

INDICATOR STANDARD

972

Recovery rate

83%

> 75%



Default rate

2%

< 15%



Death rate

0%

< 3%



Referral rate

2%

## Water, Sanitation and Hygiene (WASH)

### WATER

Average number of litres of potable water available per person per day

INDICATOR STANDARD

21

< 20



Proportion of households collecting >= 15 litres per person per day

Not collected

Number of persons of concern per water tap

Not collected

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5

Not collected

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml

Not collected

Proportion of households collecting drinking water from protected water sources only

Not collected

Proportion of households with sufficient daily water storage capacity

Not collected

### WATER STAFFING

Number of hygiene promoters

INDICATOR STANDARD

20

1 : < 20,000



### SANITATION

Refugees per latrine/toilet

INDICATOR STANDARD

15

< 20



Proportion of households with drop-hole latrine or drop-hole toilet

Not collected

Proportion of households report defecating in a toilet

Not collected

Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

Not collected