

Site report 2018 Mole CONGO (DR)



Population


15,969 refugees

23%

Children under 5

20%

Women of reproductive age



Origin of refugees

RCA

Site opened
Jun 2013

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.2

Deaths/1,000 refugees/month

MATERNAL MORTALITY

1

Deaths

SKILLED BIRTH ATTENDANCE

97%

of all births

GAM

NC

Global Acute Malnutrition

SAM

NC

Severe Acute Malnutrition

WATER SUPPLY

21


Liters/ person/ day

LATRINE COVERAGE



22

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING

| | INDICATOR | STANDARD |
|------------------------------------|-----------|--------------|
| Number of medical doctors | 1 | 1 : < 50,000 |
| Number of qualified nurses | 12 | 1 : < 10,000 |
| Number of community health workers | 15 | 1 : < 1,000 |

ACCESS AND UTILIZATION

| | INDICATOR | STANDARD |
|---|-----------|----------|
| Consultations per trained clinician per day | 17 | < 50 |
| Health utilisation rate (new visits / refugee / year) | 0.9 | 1 - 4 |
| Proportion of host population consultations | 15% | |

MORTALITY

| | INDICATOR | STANDARD |
|---|-----------|----------|
| Crude Mortality Rate (CMR) (/1000/month) | 0.1 | < 0.75 |
| Under-five Mortality Rate (U5MR) (/1000/month) | 0.2 | < 1.5 |
| Infant Mortality Rate (IMR) (/1000 livebirths) | 21.6 | < 30 |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths) | 0 | < 20 |

OUTBREAK ALERT AND RESPONSE

| | INDICATOR | STANDARD |
|--|-----------|----------|
| Proportion of outbreaks investigated within 48 hours | | 100% |
| Number of outbreaks reported | 0 | |

MORBIDITY

| | INDICATOR | STANDARD |
|--|-----------|----------|
| Incidence of malaria among children under 5 | 48 | |
| Incidence of watery diarrhoea among children under 5 | 0 | |
| Incidence of pneumonia among children under 5 | 1 | |
| Incidence of pneumonia among over 5 | 1 | |
| Tuberculosis success rate | 80% | < 90% |
| Were any MDR/X-TB cases diagnosed among PoCs? | No | No |

VACCINATION

| | INDICATOR | STANDARD |
|------------------------------|-----------|----------|
| Full vaccination coverage | 97% | < 95% |
| Measles vaccination coverage | 97% | < 95% |
| Source of vaccination data | HIS | |

IN-PATIENT DEPARTMENT (IPD)

| | INDICATOR | STANDARD |
|-------------------------------|-----------|----------|
| Average length of stay (days) | 0.7 | |
| Case fatality rate | 0.0 | |
| Hospitalisation rate | 53.9 | 50 - 150 |

Reproductive Health

ANTENATAL CARE

| | INDICATOR | STANDARD |
|---|-----------|----------|
| Antenatal care coverage | 80% | > 90% |
| Coverage of antenatal tetanus vaccination | 89% | > 95% |

DELIVERY CARE

| | INDICATOR | STANDARD |
|--|-----------|----------|
| Proportion of births attended by skilled personnel | 97% | ≥ 90% |
| Proportion of births conducted by caesarean section | 5% | 5 - 15% |
| Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs) | 5% | < 15% |
| Proportion of teenage pregnancies | 7% | |
| Still birth rate (/ 1000 total births / month) | 1 | |
| Number of maternal deaths | 1 | |
| Proportion of maternal deaths investigated within 48 hours | 100% | 100% |

FAMILY PLANNING

| | INDICATOR | STANDARD |
|---|-----------|----------|
| Contraceptive prevalence rate | 15% | ≥ 30% |
| Source of contraceptive prevalence data | HIS | |

SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

| | INDICATOR | STANDARD |
|--|-----------|----------|
| Total number of reported rapes | 16 | |
| Proportion of eligible rape survivors provided with PEP within 72 hours | 100% | 100% |
| Proportion of eligible rape survivors provided with ECP within 120 hours | 33% | 100% |

SEXUALLY TRANSMITTED INFECTIONS (STIS)

| | INDICATOR | STANDARD |
|--|-----------|----------|
| Number of genital ulcer diseases | 173 | |
| Number of Pelvic Inflammatory diseases | 155 | |

HIV/AIDS

PREVENTION & TREATMENT

| | INDICATOR | STANDARD |
|--|-----------|----------|
| Proportion of blood units screened for HIV | 100% | 100% |
| Condom distribution rate | 0.48 | > 0.5 |
| Proportion of TB patients tested for HIV | | 100% |

PREVENTION & TREATMENT

| | INDICATOR | STANDARD | |
|--|-----------|----------|---|
| Number of PoCs receiving ART | 122 | | ✓ |
| Proportion of HIV positive mothers who swallowed ARV during delivery | | 100% | ⚠ |
| PMTCT coverage | 100% | 100% | ✓ |

Nutrition

NUTRITION GENERAL

| | INDICATOR | STANDARD |
|--|---------------|----------|
| Prevalence of global acute malnutrition (6-59 months) | Not collected | |
| Prevalence of stunting (6-59 months) | Not collected | |
| Prevalence of severe acute malnutrition (6 - 59 months) | Not collected | |
| Prevalence of anaemia in children (6-59 months) | Not collected | |
| Prevalence of anaemia in women of reproductive age (15-49 yrs) | Not collected | |

INFANT AND YOUNG CHILD FEEDING (IYCF)

| | INDICATOR | STANDARD |
|---|---------------|----------|
| Proportion of children 0-23 months who were put to the breast within one hour of birth | Not collected | |
| Proportion of infants aged between 0-5 months who received only breast milk during the previous day | Not collected | |
| IYCF programmes targeting pregnant and lactating women established or maintained? | | Yes |

MICRONUTRIENT DEFICIENCY

| | INDICATOR | STANDARD |
|---|---------------|----------|
| Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF) | Not collected | |

CMAM: SAM TREATMENT

| | INDICATOR | STANDARD |
|---|-----------|----------|
| Number of new severe acute admissions (6-59 months) | 50 | |
| Recovery rate | 100% | > 75% |
| Default rate | 0% | < 0% |
| Death rate | 0% | < 10% |
| Referral rate | 0% | |

MAM TREATMENT

| | INDICATOR | STANDARD | |
|---|-----------|----------|---|
| Number of new moderate acute admissions (6-59 months) | 36 | | |
| Recovery rate | 85% | > 75% | ✓ |
| Default rate | 0% | < 15% | ✓ |
| Death rate | 0% | < 3% | ✓ |
| Referral rate | 15% | | |

Water, Sanitation and Hygiene (WASH)

WATER

| | INDICATOR | STANDARD | |
|--|-----------|----------|---|
| Average number of litres of potable water available per person per day | 21 | < 20 | ✓ |
| Proportion of households collecting >= 15 litres per person per day | 51% | > 95% | ✗ |
| Number of persons of concern per water tap | 523 | < 80 | ✗ |
| Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5 | 100% | > 95% | ✓ |
| Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml | 100% | > 95% | ✓ |
| Proportion of households collecting drinking water from protected water sources only | 99% | > 95% | ✓ |
| Proportion of households with sufficient daily water storage capacity | 63% | > 80% | ⚠ |

WATER STAFFING

| | INDICATOR | STANDARD | |
|-----------------------------|-----------|--------------|---|
| Number of hygiene promoters | 1 | 1 : < 20,000 | ✓ |

| | INDICATOR | STANDARD | |
|---|-----------|----------|---|
| Refugees per latrine/toilet | 22 | < 20 | ⚠ |
| Proportion of households with drop-hole latrine or drop-hole toilet | 63% | | |
| Proportion of households report defecating in a toilet | 32% | > 85% | ✗ |
| Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety) | 90% | > 90% | ✓ |

SANITATION