

Site report 2018 Batouri CAMEROON



Population


5,222 refugees

16%

Children under 5

20%

Women of reproductive age



Origin of refugees

CAMEROON

Site opened
Oct 2014

Partners



Key Indicators

CRUDE MORTALITY RATE

1.2

Deaths/1,000 refugees/month

U5 MORTALITY RATE

3.6

Deaths/1,000 refugees/month

MATERNAL MORTALITY

2

Deaths

SKILLED BIRTH ATTENDANCE

84%

of all births

GAM

NC

Global Acute Malnutrition

SAM

NC

Severe Acute Malnutrition


WATER SUPPLY

Liters/ person/ day


LATRINE COVERAGE

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000	⊖	Incidence of malaria among children under 5	147
Number of qualified nurses		1 : < 10,000	⊖	Incidence of watery diarrhoea among children under 5	23
Number of community health workers		1 : < 1,000	⊖	Incidence of pneumonia among children under 5	11
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	9	< 50	✓	Incidence of pneumonia among over 5	4
Health utilisation rate (new visits / refugee / year)	1.4	1 - 4	✓	Tuberculosis success rate	< 90%
Proportion of host population consultations	43%			Were any MDR/X-TB cases diagnosed among PoCs?	No
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	1.2	< 0.75	✗	Full vaccination coverage	99% < 95% ✓
Under-five Mortality Rate (U5MR) (/1000/month)	3.6	< 1.5	✗	Measles vaccination coverage	99% < 95% ✓
Infant Mortality Rate (IMR) (/1000 livebirths)	102.7	< 30	✗	Source of vaccination data	
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	59	< 20	✗	Average length of stay (days)	11.9
OUTBREAK ALERT AND RESPONSE			CASE FATALITY RATE		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0
Number of outbreaks reported	0			Hospitalisation rate	346.0 50 - 150 ✓

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	34%	> 90%	✗	Contraceptive prevalence rate	8% ≥ 30% ✗
Coverage of antenatal tetanus vaccination	50%	> 95%	✗	Source of contraceptive prevalence data	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	84%	≥ 90%	⚠	Total number of reported rapes	0
Proportion of births conducted by caesarean section	10%	5 - 15%	✓	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	13%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%
Proportion of teenage pregnancies	15%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	5			Number of genital ulcer diseases	33
Number of maternal deaths	2			Number of Pelvic Inflammatory diseases	146
Proportion of maternal deaths investigated within 48 hours	0%	100%	✗		

HIV/AIDS

PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%
Condom distribution rate	0.00	> 0.5
Proportion of TB patients tested for HIV		100%

PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	41		✓
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✗
PMTCT coverage	120%	100%	✓

Nutrition

NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	Not collected	
Prevalence of stunting (6-59 months)	Not collected	
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected	
Prevalence of anaemia in children (6-59 months)	Not collected	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected	

INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD
Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected	
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected	
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes

MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%

CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	70	
Recovery rate	74%	> 75%
Default rate	5%	< 0%
Death rate	0%	< 10%
Referral rate	19%	

MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)			
Recovery rate		> 75%	⚠
Default rate		< 15%	✗
Death rate		< 3%	✓
Referral rate			

Water, Sanitation and Hygiene (WASH)

WATER

	INDICATOR	STANDARD
Average number of litres of potable water available per person per day		< 20
Proportion of households collecting >= 15 litres per person per day		> 95%
Number of persons of concern per water tap		< 80
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%
Proportion of households collecting drinking water from protected water sources only		> 95%
Proportion of households with sufficient daily water storage capacity		> 80%

WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters	1 : < 20,000		⊖

SANITATION

	INDICATOR	STANDARD
Refugees per latrine/toilet		< 20
Proportion of households with drop-hole latrine or drop-hole toilet		
Proportion of households report defecating in a toilet		> 85%
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%