

# Site report 2018 Belita CAMEROON



## Population


**1,303** refugees

**19%**

Children under 5

**20%**

Women of reproductive age



Origin of refugees

**NOT COLLECTED**

Site opened

**Invalid date**

## Partners

## Key Indicators

CRUDE MORTALITY RATE

**0.1**

Deaths/1,000 refugees/month

U5 MORTALITY RATE

**0.3**

Deaths/1,000 refugees/month

MATERNAL MORTALITY

**0**

Deaths

SKILLED BIRTH ATTENDANCE

**59%**

of all births

GAM

**NC**

Global Acute Malnutrition

SAM

**NC**

Severe Acute Malnutrition


WATER SUPPLY

Liters/ person/ day


LATRINE COVERAGE

Refugees/communal toilets or latrines


### COMMUNICABLE DISEASES




### INJURIES



### NON-COMMUNICABLE DISEASES



### MENTAL HEALTH



## Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000	⊖	Incidence of malaria among children under 5	210
Number of qualified nurses		1 : < 10,000	⊖	Incidence of watery diarrhoea among children under 5	10
Number of community health workers		1 : < 1,000	⊖	Incidence of pneumonia among children under 5	56
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	15	< 50	✓	Incidence of pneumonia among over 5	15
Health utilisation rate (new visits / refugee / year)	2.9	1 - 4	✓	Tuberculosis success rate	< 90%
Proportion of host population consultations	22%			Were any MDR/X-TB cases diagnosed among PoCs?	No
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	✓	Full vaccination coverage	171% < 95% ✓
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5	✓	Measles vaccination coverage	206% < 95% ✓
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	✓	Source of vaccination data	
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	✓	Average length of stay (days)	2.6
OUTBREAK ALERT AND RESPONSE			Case fatality rate		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0
Number of outbreaks reported	0			Hospitalisation rate	178.8 50 - 150 ✓

## Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	16%	> 90%	✗	Contraceptive prevalence rate	0% ≥ 30% ✗
Coverage of antenatal tetanus vaccination	44%	> 95%	✗	Source of contraceptive prevalence data	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	59%	≥ 90%	✗	Total number of reported rapes	0
Proportion of births conducted by caesarean section	0%	5 - 15%	✗	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	6%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%
Proportion of teenage pregnancies	9%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate ( / 1000 total births / month)	0			Number of genital ulcer diseases	12
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	28
Proportion of maternal deaths investigated within 48 hours		100%			

## HIV/AIDS

### PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.00	> 0.5
Proportion of TB patients tested for HIV		100%

### PREVENTION & TREATMENT

	INDICATOR	STANDARD
Number of PoCs receiving ART	1	
Proportion of HIV positive mothers who swallowed ARV during delivery	✗	100%
PMTCT coverage	14%	100%

## Nutrition

### NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	Not collected	
Prevalence of stunting (6-59 months)	Not collected	
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected	
Prevalence of anaemia in children (6-59 months)	Not collected	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected	

### INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD
Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected	
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected	
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes

### MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%

### CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	59	
Recovery rate	76%	> 75%
Default rate	2%	< 0%
Death rate	0%	< 10%
Referral rate	22%	

### MAM TREATMENT

	INDICATOR	STANDARD
Number of new moderate acute admissions (6-59 months)		
Recovery rate	✓	> 75%
Default rate	✗	< 15%
Death rate	✓	< 3%
Referral rate		

## Water, Sanitation and Hygiene (WASH)

### WATER

	INDICATOR	STANDARD
Average number of litres of potable water available per person per day		< 20
Proportion of households collecting >= 15 litres per person per day		> 95%
Number of persons of concern per water tap		< 80
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5		> 95%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%
Proportion of households collecting drinking water from protected water sources only		> 95%
Proportion of households with sufficient daily water storage capacity		> 80%

### WATER STAFFING

	INDICATOR	STANDARD
Number of hygiene promoters		1 : < 20,000

### SANITATION

	INDICATOR	STANDARD
Refugees per latrine/toilet		< 20
Proportion of households with drop-hole latrine or drop-hole toilet		
Proportion of households report defecating in a toilet		> 85%
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%