
Site report 2018 Garga Sarali CAMEROON

Population

5,800 refugees





Refugee Agency

Origin of refugees NOT COLLECTED

Site opened **Invalid date**

Partners

Key Indicators

CRUDE MORTALITY RATE

0.0

Deaths/1,000 refugees/

GAM

NC

Global Acute Malnutrition **U5 MORTALITY RATE**

Deaths/1,000 refugees/

SAM

NC

Severe Acute Malnutrition

MATERNAL MORTALITY

Deaths

WATER SUPPLY

Liters/ person/ day


SKILLED BIRTH **ATTENDANCE**


68% of all births

LATRINE COVERAGE


Refugees/communal toilets or latrines

COMMUNICABLE DISEASES





NON-COMMUNICABLE DISEASES



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STA	ANDARD	
Number of medical doctors		1:<50,000	•	Incidence of malaria among children under 5	17			
Number of qualified nurses		1:<10,000	•	Incidence of watery diarrhoea among	1			
Number of community health workers		1 : < 1,000	•	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	3			
Consultations per trained clinician per day	11	< 50	•	Incidence of pneumonia among over 5	1			
Health utilisation rate (new visits / refugee /	0.3	1 - 4	8	Tuberculosis success rate			< 90%	
year)			Were any MDR/X-TB cases diagnosed among PoCs?		No			
Proportion of host population consultations	66%			rucs:				
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICAT	OR	STANDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	②	Full vaccination coverage	4%		< 95%	•
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5	②	Measles vaccination coverage	4%		< 95%	•
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	②	Source of vaccination data				
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICAT	OR	STANDARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	17.7			
Proportion of outbreaks investigated within 48 hours	100%	100%	②	Case fatality rate	0.0			
Number of outbreaks reported	6			Hospitalisation rate	28.3		50 - 150	•

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	3%	> 90%	8	Contraceptive prevalence rate	0%	≥ 30%	€
Coverage of antenatal tetanus vaccination	14%	> 95%	8	Source of contraceptive prevalence data			
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	68%	≥ 90%	8	Total number of reported rapes	1		
Proportion of births conducted by caesarean section	6%	5 - 15%	②	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	6%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours	0%	100%	•
Proportion of teenage pregnancies	15%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	2			Number of genital ulcer diseases	0		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0		
Proportion of maternal deaths investigated within 48 hours	100%	100%	②				

HIV/AIDS

Death rate

Referral rate

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	②	Number of PoCs receving ART	0	
Condom distribution rate	0.00	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	72%	100%
Nutrition						
NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	Not	collected		Proportion of children 0-23 months who were put to the breast within one hour of birth	Not	collected
Prevalence of stunting (6-59 months)	Not	collected		Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not	collected
Prevalence of severe acute malnutrition (6 - 59 months)	Not	collected		IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	Not	collected		MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not	collected		Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	44			Number of new moderate acute admissions (6-59 months)	0	
Recovery rate	98%	> 75%	Ø	Recovery rate		> 75%
Default rate	0%	< 0%	8	Default rate		< 15%

Water, Sanitation and Hygiene (WASH)

0%

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day		< 20	Number of hygiene promoters		1:<20,000
Proportion of households collecting >= 15 litres per person per day		> 95%	SANITATION Refugees per latrine/toilet	INDICATOR	STANDARD < 20
Number of persons of concern per water tap		< 80	Proportion of households with drop-hole		20
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%	latrine or drop-hole toilet Proportion of households report defecating in a toilet		> 85%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%
Proportion of households collecting drinking water from protected water sources only		> 95%			
Proportion of households with sufficient daily water storage capacity		> 80%			

< 10%

Death rate

Referral rate

< 3%