

33.3 %

weapon)

#### Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD	
Number of medical doctors		1 : < 50,000	•	Incidence of malaria among children under 5	107		
Number of qualified nurses		1 : < 10,000	•	Incidence of watery diarrhoea among	3		
Number of community health workers		1 : < 1,000	•	children under 5			
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	1		
Consultations per trained clinician per day	19	< 50	0	Incidence of pneumonia among over 5	1		
Health utilisation rate (new visits / refugee /	1.0	1 - 4	0	Tuberculosis success rate		< 90%	
year)				Were any MDR/X-TB cases diagnosed among		No	
Proportion of host population consultations	20%			PoCs?			
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	STANDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.7	< 0.75	0	Full vaccination coverage	44%	< 95%	8
Under-five Mortality Rate (U5MR) (/1000/month)	2.0	< 1.5	⊗	Measles vaccination coverage	44%	< 95%	8
Infant Mortality Rate (IMR) (/1000 livebirths)	125.0	< 30	8	Source of vaccination data			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	125	< 20	8	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STANDARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	5.4		
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0		
Number of outbreaks reported	0			Hospitalisation rate	210.8	50 - 150	

### Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	56%	> 90%	8	Contraceptive prevalence rate	12%	≥ 30%	8
Coverage of antenatal tetanus vaccination	55%	> 95%	8	Source of contraceptive prevalence data			
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	100%	≥ 90%	0	Total number of reported rapes	0		
Proportion of births conducted by caesarean section	9%	5 - 15%	0	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	26%	< 15%	0	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	8%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate ( / 1000 total births / month)	14			Number of genital ulcer diseases	1		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	23		
Proportion of maternal deaths investigated within 48 hours	100%	100%	0				

#### HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD	
Proportion of blood units screened for HIV	100%	100%	0	Number of PoCs receving ART	0		
Condom distribution rate	0.00	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%	
Proportion of TB patients tested for HIV		100%		PMTCT coverage	185%	100%	<b>⊘</b>

# Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not	collected
Prevalence of stunting (6-59 months)	Not	collected		Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not	collected
Prevalence of severe acute malnutrition (6 - 59 months)	Not	collected		IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	Not	collected		MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not	Not collected		Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	276			Number of new moderate acute admissions (6-59 months)	0	
Recovery rate	97%	> 75%	0	Recovery rate		> 75%
Default rate	0%	< 0%	8	Default rate		< 15%
Death rate	2%	< 10%	0	Death rate		< 3%
Referral rate	1%			Referral rate		

## Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day		< 20	Number of hygiene promoters		1:<20,000
Proportion of households collecting >= 15		> 95%	SANITATION	INDICATOR	STANDARD
litres per person per day		5577	Refugees per latrine/toilet		< 20
Number of persons of concern per water tap		< 80	Proportion of households with drop-hole latrine or drop-hole toilet		
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU		> 95%	Proportion of households report defecating		> 85%
<5					
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%
Proportion of households collecting drinking water from protected water sources only		> 95%			
Proportion of households with sufficient daily water storage capacity		> 80%			