
Site report 2018 Mborgop CAMEROON

Population

11,852 refugees







Origin of refugees

CENTRAL AFRICAN REPUBLIC



Site opened Mar 2014

Partners







Key Indicators

CRUDE MORTALITY RATE

Deaths/1,000 refugees/

GAM

NC Global Acute

U5 MORTALITY RATE

Deaths/1,000 refugees/

SAM

NC

Severe Acute Malnutrition

MATERNAL MORTALITY

Deaths

WATER SUPPLY

16.1

Liters/ person/ day


SKILLED BIRTH **ATTENDANCE**


93% of all births


LATRINE COVERAGE

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES








NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STA	NDARD	
Number of medical doctors		1:<50,000	•	Incidence of malaria among children under 5	22			
Number of qualified nurses		1 : < 10,000	•	Incidence of watery diarrhoea among	7			
Number of community health workers		1 : < 1,000	•	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	1			
Consultations per trained clinician per day	16	< 50	•	Incidence of pneumonia among over 5	0			
Health utilisation rate (new visits / refugee /	0.7	1 - 4	8	Tuberculosis success rate		<	90%	
year)	2%			Were any MDR/X-TB cases diagnosed among PoCs?	No		No	
Proportion of host population consultations	290							
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	OR	STANDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	②	Full vaccination coverage	163%		< 95%	•
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5	②	Measles vaccination coverage	80%		< 95%	8
Infant Mortality Rate (IMR) (/1000 livebirths)	4.5	< 30	⊘	Source of vaccination data				
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	5	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATO	OR	STANDARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	2.5			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	40.2		50 - 150	<u> </u>

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	62%	> 90%	8	Contraceptive prevalence rate	0%	≥ 30%	•
Coverage of antenatal tetanus vaccination	56%	> 95%	8	Source of contraceptive prevalence data			
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	93%	≥ 90%	②	Total number of reported rapes	0		
Proportion of births conducted by caesarean section	0%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	2%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	6%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	3		
Number of maternal deaths	1			Number of Pelvic Inflammatory diseases	23		
Proportion of maternal deaths investigated within 48 hours	200%	100%	②				

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	0	
Condom distribution rate	0.00	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	38%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	Not collected		Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected	
Prevalence of stunting (6-59 months)	Not collected		Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected	
Prevalence of severe acute malnutrition (6 - 59 months)			IYCF programmes targeting pregnant and lactating women established or maintained?	Yes	
Prevalence of anaemia in children (6-59 months)	Not collected		MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected		Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%
CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	0		Number of new moderate acute admissions (6-59 months)	0	
Recovery rate		> 75%	Recovery rate		> 75%
Default rate		< 0%	Default rate		< 15%
Death rate		< 10%	Death rate		< 3%
Referral rate			Referral rate		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	16	< 20	8	Number of hygiene promoters		1:<20,000	•
Proportion of households collecting >= 15 litres per person per day		> 95%		SANITATION Refugees per latrine/toilet	INDICATOR 9	STANDARD < 20	•
Number of persons of concern per water tap		< 80		Proportion of households with drop-hole	9	\20	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU		> 95%		latrine or drop-hole toilet Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%		Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	
Proportion of households collecting drinking water from protected water sources only		> 95%					
Proportion of households with sufficient daily water storage capacity		> 80%					