# Site report 2018 Meiganga CAMEROON

## **Population**

11,600 refugees







**Origin of refugees NOT COLLECTED** 

Site opened **Invalid date** 

### **Partners**

### **Key Indicators**

CRUDE MORTALITY RATE

0.0

Deaths/1,000 refugees/

**GAM** 

NC

Global Acute Malnutrition **U5 MORTALITY RATE** 

Deaths/1,000 refugees/

SAM

NC

Severe Acute Malnutrition

MATERNAL MORTALITY

Deaths

WATER SUPPLY

Liters/ person/ day


SKILLED BIRTH **ATTENDANCE** 


91% of all births

LATRINE COVERAGE


Refugees/communal toilets or latrines

#### COMMUNICABLE DISEASES





#### **NON-COMMUNICABLE DISEASES**



**MENTAL HEALTH** 

## **Public Health**

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD	
Number of medical doctors		1:<50,000	•	Incidence of malaria among children under 5	12		
Number of qualified nurses		1:<10,000	•	Incidence of watery diarrhoea among	1		
Number of community health workers		1:<1,000	•	children under 5			
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	1		
Consultations per trained clinician per day	8	< 50	•	Incidence of pneumonia among over 5	1		
Health utilisation rate (new visits / refugee / year)	0.2	1 - 4	8	Tuberculosis success rate		< 90%	
Proportion of host population consultations	65%			Were any MDR/X-TB cases diagnosed among PoCs?		No	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	STANDAR	D
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	<b>⊘</b>	Full vaccination coverage	1%	< 95%	•
Under-five Mortality Rate (U5MR) (/1000/month)	0.1	< 1.5	<b>⊘</b>	Measles vaccination coverage	7%	< 95%	•
Infant Mortality Rate (IMR) (/1000 livebirths)	23.3	< 30	<b>⊘</b>	Source of vaccination data			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	<b>②</b>	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STANDAR	D
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	11.5		
Proportion of outbreaks investigated within 48 hours	100%	100%	<b>②</b>	Case fatality rate	0.0		
Number of outbreaks reported	12			Hospitalisation rate	18.9	50 - 150	•

## **Reproductive Health**

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	30%	> 90%	8	Contraceptive prevalence rate	0%	≥ 30%	
Coverage of antenatal tetanus vaccination	30%	> 95%	8	Source of contraceptive prevalence data			
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	91%	≥ 90%	<b>②</b>	Total number of reported rapes	0		
Proportion of births conducted by caesarean section	2%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	5%	< 15%	<b>②</b>	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	14%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate ( / 1000 total births / month)	0			Number of genital ulcer diseases	0		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	7		
Proportion of maternal deaths investigated within 48 hours	100%	100%	<b>②</b>				

## HIV/AIDS

Death rate

Referral rate

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	0	
Condom distribution rate	0.00	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	91%	100%
Nutrition						
NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected	
Prevalence of stunting (6-59 months)	Not	collected		Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected	
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected			IYCF programmes targeting pregnant and lactating women established or maintained?	Yes	
Prevalence of anaemia in children (6-59 months)	Not	collected		MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD

Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected		Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	> 70%		
CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	0		Number of new moderate acute admissions (6-59 months)	0		
Recovery rate		> 75%	Recovery rate		> 75%	
Default rate		< 0%	Default rate		< 15%	

Death rate

Referral rate

< 3%

< 10%

## Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day		< 20	Number of hygiene promoters		1:<20,000	•
Proportion of households collecting >= 15 litres per person per day		> 95%	SANITATION  Refugees per latrine/toilet	INDICATOR	STANDARD < 20	
Number of persons of concern per water tap		< 80	Proportion of households with drop-hole		<b>\20</b>	
Proportion of tests at chlorinated water collection points indicating a measurable		> 95%	latrine or drop-hole toilet		> 0F0/	
chlorine residual ( >= 0.1 mg/litre) and NTU <5			Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/		> 95%	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	
		> 95%				
Proportion of households collecting drinking water from protected water sources only		~ <del>9</del> 3%0				
Proportion of households with sufficient daily water storage capacity		> 80%				