

# Site report 2018 Ngaoui CAMEROON



## Population


**1,303** refugees

**19%**

Children under 5

**20%**

Women of reproductive age



Origin of refugees

**NOT COLLECTED**

Site opened

**Invalid date**

## Partners

## Key Indicators

CRUDE MORTALITY RATE

**0.4**

Deaths/1,000 refugees/month

U5 MORTALITY RATE

**1.7**

Deaths/1,000 refugees/month

MATERNAL MORTALITY

**0**

Deaths

SKILLED BIRTH ATTENDANCE

**100%**

of all births

GAM

**NC**

Global Acute Malnutrition

SAM

**NC**

Severe Acute Malnutrition


WATER SUPPLY

Liters/ person/ day

LATRINE COVERAGE


Refugees/communal toilets or latrines

## COMMUNICABLE DISEASES



## INJURIES

## NON-COMMUNICABLE DISEASES



## MENTAL HEALTH

## Public Health

### HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000
Number of qualified nurses		1 : < 10,000
Number of community health workers		1 : < 1,000

### ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	4	< 50
Health utilisation rate (new visits / refugee / year)	0.1	1 - 4
Proportion of host population consultations	90%	

### MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.4	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	1.7	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	286	< 20

### OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours	0%	100%
Number of outbreaks reported	1	

### MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	11	
Incidence of watery diarrhoea among children under 5	0	
Incidence of pneumonia among children under 5	2	
Incidence of pneumonia among over 5	1	
Tuberculosis success rate		< 90%
Were any MDR/X-TB cases diagnosed among PoCs?		No

### VACCINATION

	INDICATOR	STANDARD
Full vaccination coverage	0%	< 95%
Measles vaccination coverage	0%	< 95%
Source of vaccination data		

### IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD
Average length of stay (days)	12.5	
Case fatality rate	0.0	
Hospitalisation rate	76.0	50 - 150

## Reproductive Health

### ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	83%	> 90%
Coverage of antenatal tetanus vaccination	150%	> 95%

### DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	100%	≥ 90%
Proportion of births conducted by caesarean section	33%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	0%	< 15%
Proportion of teenage pregnancies	0%	
Still birth rate ( / 1000 total births / month)	0	
Number of maternal deaths	0	
Proportion of maternal deaths investigated within 48 hours		100%

### FAMILY PLANNING

	INDICATOR	STANDARD
Contraceptive prevalence rate	0%	≥ 30%
Source of contraceptive prevalence data		

### SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD
Total number of reported rapes	0	
Proportion of eligible rape survivors provided with PEP within 72 hours		100%
Proportion of eligible rape survivors provided with ECP within 120 hours		100%

### SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD
Number of genital ulcer diseases	0	
Number of Pelvic Inflammatory diseases	0	

## HIV/AIDS

### PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.00	> 0.5
Proportion of TB patients tested for HIV		100%

### PREVENTION & TREATMENT

	INDICATOR	STANDARD
Number of PoCs receiving ART	0	
Proportion of HIV positive mothers who swallowed ARV during delivery	✘	100%
PMTCT coverage		100%

## Nutrition

### NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	Not collected	
Prevalence of stunting (6-59 months)	Not collected	
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected	
Prevalence of anaemia in children (6-59 months)	Not collected	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected	

### INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD
Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected	
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected	
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes

### MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%

### CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)		
Recovery rate		> 75%
Default rate		< 0%
Death rate		< 10%
Referral rate		

### MAM TREATMENT

	INDICATOR	STANDARD
Number of new moderate acute admissions (6-59 months)	0	
Recovery rate		> 75%
Default rate		< 15%
Death rate		< 3%
Referral rate		

## Water, Sanitation and Hygiene (WASH)

### WATER

	INDICATOR	STANDARD
Average number of litres of potable water available per person per day		< 20
Proportion of households collecting >= 15 litres per person per day		> 95%
Number of persons of concern per water tap		< 80
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5		> 95%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%
Proportion of households collecting drinking water from protected water sources only		> 95%
Proportion of households with sufficient daily water storage capacity		> 80%

### WATER STAFFING

	INDICATOR	STANDARD
Number of hygiene promoters		1 : < 20,000

### SANITATION

	INDICATOR	STANDARD
Refugees per latrine/toilet		< 20
Proportion of households with drop-hole latrine or drop-hole toilet		
Proportion of households report defecating in a toilet		> 85%
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%