

Site report 2018 Tongo Gandima CAMEROON



Population

3,204 refugees

16%

Children under 5

20%

Women of reproductive age



Origin of refugees

NOT COLLECTED

Site opened

Invalid date

Partners

Key Indicators

CRUDE MORTALITY RATE

0.1

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.2

Deaths/1,000 refugees/month

MATERNAL MORTALITY

0

Deaths

SKILLED BIRTH ATTENDANCE

67%

of all births

GAM

NC

Global Acute Malnutrition

SAM

NC

Severe Acute Malnutrition


WATER SUPPLY

Liters/ person/ day


LATRINE COVERAGE

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000
Number of qualified nurses		1 : < 10,000
Number of community health workers		1 : < 1,000

ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	4	< 50
Health utilisation rate (new visits / refugee / year)	0.6	1 - 4
Proportion of host population consultations	44%	

MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20

OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%
Number of outbreaks reported	0	

MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	72	
Incidence of watery diarrhoea among children under 5	38	
Incidence of pneumonia among children under 5	30	
Incidence of pneumonia among over 5	8	
Tuberculosis success rate		< 90%
Were any MDR/X-TB cases diagnosed among PoCs?		No

VACCINATION

	INDICATOR	STANDARD
Full vaccination coverage	8%	< 95%
Measles vaccination coverage	38%	< 95%
Source of vaccination data		

IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD
Average length of stay (days)	6.5	
Case fatality rate	0.0	
Hospitalisation rate	47.4	50 - 150

Reproductive Health

ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	21%	> 90%
Coverage of antenatal tetanus vaccination	41%	> 95%

DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	67%	≥ 90%
Proportion of births conducted by caesarean section	0%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	3%	< 15%
Proportion of teenage pregnancies	9%	
Still birth rate (/ 1000 total births / month)	0	
Number of maternal deaths	0	
Proportion of maternal deaths investigated within 48 hours		100%

FAMILY PLANNING

	INDICATOR	STANDARD
Contraceptive prevalence rate	3%	≥ 30%
Source of contraceptive prevalence data		

SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD
Total number of reported rapes	0	
Proportion of eligible rape survivors provided with PEP within 72 hours		100%
Proportion of eligible rape survivors provided with ECP within 120 hours		100%

SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD
Number of genital ulcer diseases	1	
Number of Pelvic Inflammatory diseases	41	

HIV/AIDS

PREVENTION & TREATMENT

INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%
Condom distribution rate	0.00 > 0.5
Proportion of TB patients tested for HIV	100%

PREVENTION & TREATMENT

INDICATOR	STANDARD
Number of PoCs receiving ART	8
Proportion of HIV positive mothers who swallowed ARV during delivery	100%
PMTCT coverage	114% 100%

Nutrition

NUTRITION GENERAL

INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	Not collected
Prevalence of stunting (6-59 months)	Not collected
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected
Prevalence of anaemia in children (6-59 months)	Not collected
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected

INFANT AND YOUNG CHILD FEEDING (IYCF)

INDICATOR	STANDARD
Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected
IYCF programmes targeting pregnant and lactating women established or maintained?	Yes

MICRONUTRIENT DEFICIENCY

INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	> 70%

CMAM: SAM TREATMENT

INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	45
Recovery rate	80% > 75%
Default rate	17% < 0%
Death rate	0% < 10%
Referral rate	2%

MAM TREATMENT

INDICATOR	STANDARD
Number of new moderate acute admissions (6-59 months)	0
Recovery rate	> 75%
Default rate	< 15%
Death rate	< 3%
Referral rate	

Water, Sanitation and Hygiene (WASH)

WATER

INDICATOR	STANDARD
Average number of litres of potable water available per person per day	< 20
Proportion of households collecting >= 15 litres per person per day	> 95%
Number of persons of concern per water tap	< 80
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	> 95%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	> 95%
Proportion of households collecting drinking water from protected water sources only	> 95%
Proportion of households with sufficient daily water storage capacity	> 80%

WATER STAFFING

INDICATOR	STANDARD
Number of hygiene promoters	1 : < 20,000

SANITATION

INDICATOR	STANDARD
Refugees per latrine/toilet	< 20
Proportion of households with drop-hole latrine or drop-hole toilet	
Proportion of households report defecating in a toilet	> 85%
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	> 90%