
Site report 2018 Awbarre ETHIOPIA

Population

11,895 refugees





Refugee Agency

Origin of refugees SOMALIA

Site opened Jan 2007

Partners









Key Indicators

CRUDE MORTALITY RATE

0

Deaths/1,000 refugees/

GAM

Global Acute

U5 MORTALITY RATE

Deaths/1,000 refugees/

SAM

NC

Severe Acute Malnutrition

MATERNAL MORTALITY

Deaths

WATER SUPPLY

12

Liters/ person/ day


SKILLED BIRTH **ATTENDANCE**


of all births

LATRINE COVERAGE


Refugees/communal toilets or latrines


COMMUNICABLE DISEASES





NON-COMMUNICABLE DISEASES





Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD
Number of medical doctors		1:<50,000	•	Incidence of malaria among children under 5	0	
Number of qualified nurses		1 : < 10,000	•	Incidence of watery diarrhoea among	46	
Number of community health workers		1 : < 1,000	•	children under 5		
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	93	
Consultations per trained clinician per day	25	< 50	•	Incidence of pneumonia among over 5	15	
Health utilisation rate (new visits / refugee / year)	0.9	1 - 4	A	Tuberculosis success rate		< 90%
Proportion of host population consultations	17%			Were any MDR/X-TB cases diagnosed among PoCs?		No
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	②	Full vaccination coverage	110%	< 95%
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5	②	Measles vaccination coverage	111%	< 95%
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	②	Source of vaccination data		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATO	R STANDARD
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	2.1	
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0	
Number of outbreaks reported	0			Hospitalisation rate	4.6	50 - 150

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD
Antenatal care coverage	95%	> 90%	②	Contraceptive prevalence rate	19%	≥ 30%
Coverage of antenatal tetanus vaccination	94%	> 95%	A	Source of contraceptive prevalence data		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	99%	≥ 90%	②	Total number of reported rapes	0	
Proportion of births conducted by caesarean section	0%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours	NR	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	0%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours		100%
Proportion of teenage pregnancies	2%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	0	
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0	
Proportion of maternal deaths investigated within 48 hours		100%				

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receving ART	-1	
Condom distribution rate	0.20	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	103%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected		Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected		Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected		IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	Not collected		MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected		Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%	
CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	0		Number of new moderate acute admissions (6-59 months)	42		
Recovery rate		> 75%	Recovery rate	100%	> 75%	②
Default rate		< 0%	Default rate	0%	< 15%	②
Death rate		< 10%	Death rate	0%	< 3%	Ø
Referral rate			Referral rate	0%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	12	< 20	8	Number of hygiene promoters		1:<20,000	•
Proportion of households collecting >= 15 litres per person per day		> 95%		SANITATION Refugees per latrine/toilet	INDICATOR	STANDARD < 20	•
Number of persons of concern per water tap		< 80		Proportion of households with drop-hole	,	\20	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU		> 95%		Proportion of households report defecating in a toilet		> 85%	
<5 Proportion of tests at unchlorinated water		> 95%		Proportion of communal latrines compliant with UNHCR standards (cleanable slabs,		> 90%	
collection points showing 0 Faecal Coliforms/ 100 ml				privacy & structural safety)			
Proportion of households collecting drinking water from protected water sources only		> 95%					
Proportion of households with sufficient daily water storage capacity		> 80%					