



Site report 2018 Aysaita ETHIOPIA



Population

15,195 refugees



Origin of refugees

ERITREA

Site opened
Jan 2008

Partners



Key Indicators

CRUDE MORTALITY RATE

0.0 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.1 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

92% of all births

GAM

18% Global Acute Malnutrition

SAM

0 Severe Acute Malnutrition


WATER SUPPLY

16 Liters/ person/ day



LATRINE COVERAGE

36 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Number of medical doctors	1	1 : < 50,000	✓	Incidence of malaria among children under 5	3	
Number of qualified nurses	10	1 : < 10,000	✓	Incidence of watery diarrhoea among children under 5	98	
Number of community health workers	19	1 : < 1,000	✓	Incidence of pneumonia among children under 5	65	
ACCESS AND UTILIZATION			VACCINATION			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Consultations per trained clinician per day	23	< 50	✓	Incidence of pneumonia among over 5	13	
Health utilisation rate (new visits / refugee / year)	1.1	1 - 4	✓	Tuberculosis success rate	100% < 90% ✓	
Proportion of host population consultations	0%			Were any MDR/X-TB cases diagnosed among PoCs?	No No ✗	
MORTALITY			IN-PATIENT DEPARTMENT (IPD)			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	✓	Full vaccination coverage	63% < 95% ✗	
Under-five Mortality Rate (U5MR) (/1000/month)	0.1	< 1.5	✓	Measles vaccination coverage	88% < 95% ✗	
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	✓	Source of vaccination data	HIS	
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	✓	IN-PATIENT DEPARTMENT (IPD)		
OUTBREAK ALERT AND RESPONSE				INDICATOR	STANDARD	
	INDICATOR	STANDARD				
Proportion of outbreaks investigated within 48 hours		100%		Average length of stay (days)	3.1	
Number of outbreaks reported	0			Case fatality rate	0.0	
				Hospitalisation rate	25.4 50 - 150 ✗	

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Antenatal care coverage	NR			Contraceptive prevalence rate	1% ≥ 30% ✗	
Coverage of antenatal tetanus vaccination	80%	> 95%	⚠	Source of contraceptive prevalence data	HIS	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	92%	≥ 90%	✓	Total number of reported rapes	0	
Proportion of births conducted by caesarean section	1%	5 - 15%	✗	Proportion of eligible rape survivors provided with PEP within 72 hours	100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	2%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%	
Proportion of teenage pregnancies	1%			SEXUALLY TRANSMITTED INFECTIONS (STIS)		
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	0	
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0	
Proportion of maternal deaths investigated within 48 hours		100%				

HIV/AIDS

PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.10	> 0.5
Proportion of TB patients tested for HIV		100%

PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	0		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✗
PMTCT coverage	88%	100%	✗

Nutrition

NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	18%	< 20%
Prevalence of stunting (6-59 months)	49%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	0%	< 3%
Prevalence of anaemia in children (6-59 months)	40%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	29%	< 20%

INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	94%	> 80%	✗
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	90%	> 70%	✗
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	✓

MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%

CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	486	
Recovery rate	65%	> 75%
Default rate	7%	< 0%
Death rate	1%	< 10%
Referral rate	5%	

MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	549		
Recovery rate	63%	> 75%	✗
Default rate	6%	< 15%	✓
Death rate	0%	< 3%	✓
Referral rate	10%		

Water, Sanitation and Hygiene (WASH)

WATER

	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	16	< 20	✗
Proportion of households collecting >= 15 litres per person per day		> 95%	
Number of persons of concern per water tap		< 80	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	
Proportion of households collecting drinking water from protected water sources only		> 95%	
Proportion of households with sufficient daily water storage capacity		> 80%	

WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters	5	1 : < 20,000	✓

SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	36	< 20	✗
Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of households report defecating in a toilet		> 85%	
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	