

Site report 2018 Bambasi ETHIOPIA



Population


17,103 refugees

24%

Children under 5

20%

Women of reproductive age



Origin of refugees

SUDAN

SOUTH SUDAN

Site opened
Jan 2012

Partners



Key Indicators

CRUDE MORTALITY RATE

0.0

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.1

Deaths/1,000 refugees/month

MATERNAL MORTALITY

0

Deaths

SKILLED BIRTH ATTENDANCE

99%

of all births

GAM

5%

Global Acute Malnutrition

SAM

0

Severe Acute Malnutrition

WATER SUPPLY

14


Liters/ person/ day

LATRINE COVERAGE



5

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	1	1 : < 50,000	✓	0	
Number of qualified nurses	10	1 : < 10,000	✓	7	
Number of community health workers	69	1 : < 1,000	✓	24	
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	27	< 50	✓	62%	< 95%
Health utilisation rate (new visits / refugee / year)	0.8	1 - 4	✗	96%	< 95%
Proportion of host population consultations	23%			HIS	
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	✓	10.6	
Under-five Mortality Rate (U5MR) (/1000/month)	0.1	< 1.5	✓	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	5.9	< 30	✓	Hospitalisation rate	6.8
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	4	< 20	✓		50 - 150
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%			
Number of outbreaks reported	0				

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	92%	> 90%	✓	51%	≥ 30%
Coverage of antenatal tetanus vaccination	99%	> 95%	✓	HIS	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	99%	≥ 90%	✓	1	
Proportion of births conducted by caesarean section	0%	5 - 15%	✗		100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	1%	< 15%	✓		100%
Proportion of teenage pregnancies	2%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	0
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0
Proportion of maternal deaths investigated within 48 hours		100%			

HIV/AIDS

PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.15	> 0.5
Proportion of TB patients tested for HIV		100%

PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	0		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✗
PMTCT coverage	92%	100%	⚠

Nutrition

NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	5%	< 20%
Prevalence of stunting (6-59 months)	43%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	0%	< 3%
Prevalence of anaemia in children (6-59 months)	23%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	7%	< 20%

INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	94%	> 80%	✓
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	93%	> 70%	✓
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	

MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD	
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FPB)	100%	> 70%	✓

CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	40	
Recovery rate	100%	> 75%
Default rate	0%	< 0%
Death rate	0%	< 10%
Referral rate	0%	

MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	206		
Recovery rate	99%	> 75%	✓
Default rate	0%	< 15%	✓
Death rate	0%	< 3%	✓
Referral rate	1%		

Water, Sanitation and Hygiene (WASH)

WATER

	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	14	< 20	✗
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	✓
Number of persons of concern per water tap	57	< 80	✓
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%	✓
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	✗
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	✓
Proportion of households with sufficient daily water storage capacity		> 80%	

WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters	8	1 : < 20,000	✓

SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	5	< 20	✓
Proportion of households with drop-hole latrine or drop-hole toilet	6%		
Proportion of households report defecating in a toilet	78%	> 85%	⚠
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	3%	> 90%	✗