



Site report 2018 Bokolmanyo ETHIOPIA



Population

43,543 refugees



Origin of refugees

SOMALI

Site opened
Jan 2010

Partners



Key Indicators

CRUDE MORTALITY RATE

0.0 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.1 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

100% of all births

GAM

12% Global Acute Malnutrition

SAM

2% Severe Acute Malnutrition


WATER SUPPLY

16 Liters/ person/ day



LATRINE COVERAGE

43 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000
Number of qualified nurses		1 : < 10,000
Number of community health workers		1 : < 1,000

ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	55	< 50
Health utilisation rate (new visits / refugee / year)	0.8	1 - 4
Proportion of host population consultations	11%	

MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.1	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	2.1	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20

OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%
Number of outbreaks reported	0	

MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	0	
Incidence of watery diarrhoea among children under 5	25	
Incidence of pneumonia among children under 5	26	
Incidence of pneumonia among over 5	5	
Tuberculosis success rate		< 90%
Were any MDR/X-TB cases diagnosed among PoCs?		No

VACCINATION

	INDICATOR	STANDARD	
Full vaccination coverage	60%	< 95%	✗
Measles vaccination coverage	59%	< 95%	✗
Source of vaccination data			✓

IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD	
Average length of stay (days)	4.7		
Case fatality rate	0.0		
Hospitalisation rate	6.6	50 - 150	✗

Reproductive Health

ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	35%	> 90%
Coverage of antenatal tetanus vaccination	96%	> 95%

DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	100%	≥ 90%
Proportion of births conducted by caesarean section	0%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	2%	< 15%
Proportion of teenage pregnancies	4%	
Still birth rate (/ 1000 total births / month)	2	
Number of maternal deaths	0	
Proportion of maternal deaths investigated within 48 hours	0%	100%

FAMILY PLANNING

	INDICATOR	STANDARD	
Contraceptive prevalence rate	5%	≥ 30%	✗
Source of contraceptive prevalence data			✓

SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD	
Total number of reported rapes	0		✓
Proportion of eligible rape survivors provided with PEP within 72 hours		100%	✗
Proportion of eligible rape survivors provided with ECP within 120 hours		100%	✓

SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD	
Number of genital ulcer diseases	114		
Number of Pelvic Inflammatory diseases	5		



HIV/AIDS

PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.22	> 0.5
Proportion of TB patients tested for HIV		100%

PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	7		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✗
PMTCT coverage	92%	100%	⚠

Nutrition

NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	12%	< 20%
Prevalence of stunting (6-59 months)	27%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	2%	< 3%
Prevalence of anaemia in children (6-59 months)	45%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	37%	< 20%

INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	83%	> 80%	✗
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	84%	> 70%	✗
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	

MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%

CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	1173	
Recovery rate	95%	> 75%
Default rate	0%	< 0%
Death rate	0%	< 10%
Referral rate	2%	

MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	2010		
Recovery rate	93%	> 75%	✓
Default rate	1%	< 15%	✓
Death rate	0%	< 3%	✓
Referral rate	5%		

Water, Sanitation and Hygiene (WASH)

WATER

	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	16	< 20	✗
Proportion of households collecting >= 15 litres per person per day		> 95%	
Number of persons of concern per water tap		< 80	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	
Proportion of households collecting drinking water from protected water sources only		> 95%	
Proportion of households with sufficient daily water storage capacity		> 80%	

WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters		1 : < 20,000	⊖

SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	43	< 20	✗
Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of households report defecating in a toilet		> 85%	
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	