



# Site report 2018 Fugnido ETHIOPIA



## Population

**67,437** refugees



Origin of refugees  
**SOUTH SUDAN**

Site opened  
**Apr 1993**

## Partners



## Key Indicators

CRUDE MORTALITY RATE

**0.0** Deaths/1,000 refugees/month

U5 MORTALITY RATE

**0.1** Deaths/1,000 refugees/month

MATERNAL MORTALITY

**0** Deaths

SKILLED BIRTH ATTENDANCE

**99%** of all births

GAM

**16%** Global Acute Malnutrition

SAM

**3%** Severe Acute Malnutrition


WATER SUPPLY

**19** Liters/ person/ day



LATRINE COVERAGE

**10** Refugees/communal toilets or latrines


### COMMUNICABLE DISEASES




### INJURIES



### NON-COMMUNICABLE DISEASES



### MENTAL HEALTH



## Public Health

| HEALTH STAFFING                                       |           |              | MORBIDITY                               |  |            |
|---|-----------|--------------|---|--|------------|
|   | INDICATOR | STANDARD     |   | INDICATOR  | STANDARD   |
| Number of medical doctors                             | 1         | 1 : < 50,000 | ✘                                       | 4  |            |
| Number of qualified nurses                            | 27        | 1 : < 10,000 | ✔                                       | 27   |            |
| Number of community health workers                    | 125       | 1 : < 1,000  | ✔                                       | 27   |            |
| ACCESS AND UTILIZATION                                |           |              | VACCINATION                             |  |            |
|   | INDICATOR | STANDARD     |   | INDICATOR  | STANDARD   |
| Consultations per trained clinician per day           | 15        | < 50         | ✔                                       | 9  |            |
| Health utilisation rate (new visits / refugee / year) | 0.6       | 1 - 4        | ✘                                       | 90%  | < 90% ✔    |
| Proportion of host population consultations           | 8%        |              |   | No   | No ✘       |
|   |           |              |   | Were any MDR/X-TB cases diagnosed among PoCs?                            |            |
| MORTALITY   |           |              | IN-PATIENT DEPARTMENT (IPD)             |  |            |
|   | INDICATOR | STANDARD     |   | INDICATOR  | STANDARD   |
| Crude Mortality Rate (CMR) (/1000/month)              | 0.0       | < 0.75       | ✔                                       | 2.2  |            |
| Under-five Mortality Rate (U5MR) (/1000/month)        | 0.1       | < 1.5        | ✔                                       | 0.0  |            |
| Infant Mortality Rate (IMR) (/1000 livebirths)        | 0.7       | < 30         | ✔                                       | 8.3  | 50 - 150 ✘ |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths)     | 0         | < 20         | ✔                                       |  |            |
| OUTBREAK ALERT AND RESPONSE                           |           |              | SEXUAL AND GENDER-BASED VIOLENCE (SGBV) |  |            |
|   | INDICATOR | STANDARD     |   | INDICATOR  | STANDARD   |
| Proportion of outbreaks investigated within 48 hours  |           | 100%         |   | 0  |            |
| Number of outbreaks reported                          | 0         |              |   |  |            |
|   |           |              |   | Total number of reported rapes   | 0          |
|   |           |              |   | Proportion of eligible rape survivors provided with PEP within 72 hours  | 100%       |
|   |           |              |   | Proportion of eligible rape survivors provided with ECP within 120 hours | 100%       |

## Reproductive Health

| ANTENATAL CARE   |           |          | FAMILY PLANNING                        |           |          |
|--|-----------|----------|--|-----------|----------|
|  | INDICATOR | STANDARD |  | INDICATOR | STANDARD |
| Antenatal care coverage  | 75%       | > 90%    | ✘                                      | 62%       | ≥ 30% ✔  |
| Coverage of antenatal tetanus vaccination  | 79%       | > 95%    | ⚠                                      | HIS       |          |
| DELIVERY CARE  |           |          | SEXUALLY TRANSMITTED INFECTIONS (STIS) |           |          |
|  | INDICATOR | STANDARD |  | INDICATOR | STANDARD |
| Proportion of births attended by skilled personnel                                     | 99%       | ≥ 90%    | ✔                                      | 61        |          |
| Proportion of births conducted by caesarean section                                    | 0%        | 5 - 15%  | ✘                                      | 28        |          |
| Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs) | 1%        | < 15%    | ✔                                      |           |          |
| Proportion of teenage pregnancies  | 5%        |          |  |           |          |
| Still birth rate ( / 1000 total births / month)  | 1         |          |  |           |          |
| Number of maternal deaths  | 0         |          |  |           |          |
| Proportion of maternal deaths investigated within 48 hours                             |           | 100%     |  |           |          |

## HIV/AIDS

### PREVENTION & TREATMENT

Proportion of blood units screened for HIV

Condom distribution rate

Proportion of TB patients tested for HIV

### INDICATOR

0.06

### STANDARD

100%

> 0.5

100%

### PREVENTION & TREATMENT

Number of PoCs receiving ART

✗ Proportion of HIV positive mothers who swallowed ARV during delivery

PMTCT coverage

### INDICATOR

814

100%

85%

100%



## Nutrition

### NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

Prevalence of stunting (6-59 months)

Prevalence of severe acute malnutrition (6 - 59 months)

Prevalence of anaemia in children (6-59 months)

Prevalence of anaemia in women of reproductive age (15-49 yrs)

### INDICATOR

16%

4%

3%

36%

22%

### STANDARD

< 20%

< 20%

< 3%

< 20%

< 20%

### INFANT AND YOUNG CHILD FEEDING (IYCF)

⚠ Proportion of children 0-23 months who were put to the breast within one hour of birth

✔ Proportion of infants aged between 0-5 months who received only breast milk during the previous day

⚠ IYCF programmes targeting pregnant and lactating women established or maintained?

### INDICATOR

95%

96%

Yes

### INDICATOR

100%

### STANDARD

> 80%

> 70%

Yes

### STANDARD

> 70%



### CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

Recovery rate

Default rate

Death rate

Referral rate

### INDICATOR

543

81%

3%

0%

0%

### STANDARD

> 75%

< 0%

< 10%

< 10%

### MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

✔ Recovery rate

✗ Default rate

✔ Death rate

Referral rate

### INDICATOR

1944

84%

4%

0%

9%

### STANDARD

> 75%

< 15%

< 3%



## Water, Sanitation and Hygiene (WASH)

### WATER

Average number of litres of potable water available per person per day

Proportion of households collecting >= 15 litres per person per day

Number of persons of concern per water tap

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/100 ml

Proportion of households collecting drinking water from protected water sources only

Proportion of households with sufficient daily water storage capacity

### INDICATOR

19

100%

60

100%

0%

100%

80%

### STANDARD

< 20

> 95%

< 80

> 95%

> 95%

> 95%

> 80%



### WATER STAFFING

Number of hygiene promoters

### SANITATION

Refugees per latrine/toilet

✔ Proportion of households with drop-hole latrine or drop-hole toilet

✔ Proportion of households report defecating in a toilet

✗ Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

### INDICATOR

54

### INDICATOR

10

44%

95%

100%

### STANDARD

1 : < 20,000

< 20

> 85%

> 90%

