



Site report 2018 Hilaweyn ETHIOPIA



Population

51,566 refugees



Origin of refugees

SOMALIA

Site opened
Aug 2011

Partners



Key Indicators

CRUDE MORTALITY RATE

0.0 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.3 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

95% of all births

GAM

10% Global Acute Malnutrition

SAM

2% Severe Acute Malnutrition


WATER SUPPLY

15 Liters/ person/ day



LATRINE COVERAGE

63 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000	⊖	Incidence of malaria among children under 5	0
Number of qualified nurses		1 : < 10,000	⊖	Incidence of watery diarrhoea among children under 5	36
Number of community health workers		1 : < 1,000	⊖	Incidence of pneumonia among children under 5	34
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	24	< 50	✓	Incidence of pneumonia among over 5	4
Health utilisation rate (new visits / refugee / year)	0.5	1 - 4	✗	Tuberculosis success rate	< 90%
Proportion of host population consultations	7%			Were any MDR/X-TB cases diagnosed among PoCs?	No
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	✓	Average length of stay (days)	7.1
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5	✓	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	13.3	< 30	✓	Hospitalisation rate	4.8
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	4	< 20	✓		
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%			
Number of outbreaks reported	0				

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	57%	> 90%	✗	Contraceptive prevalence rate	0%
Coverage of antenatal tetanus vaccination	94%	> 95%	⚠	Source of contraceptive prevalence data	≥ 30%
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	95%	≥ 90%	✓	Total number of reported rapes	1
Proportion of births conducted by caesarean section	0%	5 - 15%	✗	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	1%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%
Proportion of teenage pregnancies	3%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	0
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	1
Proportion of maternal deaths investigated within 48 hours		100%			

HIV/AIDS

PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.00	> 0.5
Proportion of TB patients tested for HIV		100%

PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	1		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✗
PMTCT coverage	100%	100%	⚠

Nutrition

NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	10%	< 20%
Prevalence of stunting (6-59 months)	50%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	2%	< 3%
Prevalence of anaemia in children (6-59 months)	45%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	30%	< 20%

INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	78%	> 80%	✗
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	65%	> 70%	✗
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	✓

MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/BBF)		> 70%

CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	592	
Recovery rate	96%	> 75%
Default rate	2%	< 0%
Death rate	1%	< 10%
Referral rate	1%	

MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	1512		
Recovery rate	97%	> 75%	✓
Default rate	0%	< 15%	✓
Death rate	0%	< 3%	✓
Referral rate	3%		

Water, Sanitation and Hygiene (WASH)

WATER

	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	15	< 20
Proportion of households collecting >= 15 litres per person per day		> 95%
Number of persons of concern per water tap		< 80
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%
Proportion of households collecting drinking water from protected water sources only		> 95%
Proportion of households with sufficient daily water storage capacity		> 80%

WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters		1 : < 20,000	⊖

SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	63	< 20	✗
Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of households report defecating in a toilet		> 85%	
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	