



# Site report 2018 Hitsats ETHIOPIA



## Population

**79,109** refugees



Origin of refugees

**ERITREA**

Site opened

**May 2013**

## Partners



## Key Indicators

CRUDE MORTALITY RATE

**0** Deaths/1,000 refugees/month

U5 MORTALITY RATE

**0** Deaths/1,000 refugees/month

MATERNAL MORTALITY

**0** Deaths

SKILLED BIRTH ATTENDANCE

**85%** of all births

GAM

**4%** Global Acute Malnutrition

SAM

**0%** Severe Acute Malnutrition


WATER SUPPLY

**15** Liters/ person/ day



LATRINE COVERAGE

**73** Refugees/communal toilets or latrines


### COMMUNICABLE DISEASES




### INJURIES



### NON-COMMUNICABLE DISEASES



### MENTAL HEALTH



## Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	1	1 : < 50,000	✘	36	
Number of qualified nurses	10	1 : < 10,000	✔	27	
Number of community health workers	38	1 : < 1,000	✘	44	
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	58	< 50	⚠	43%	< 95%
Health utilisation rate (new visits / refugee / year)	0.4	1 - 4	✘	97%	< 95%
Proportion of host population consultations	39%			HIS	
				100%	< 90%
				No	No
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	✔	Average length of stay (days)	7.9
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5	✔	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	✔	Hospitalisation rate	10.1
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	✔		50 - 150
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD			
Proportion of outbreaks investigated within 48 hours		100%			
Number of outbreaks reported	0				

## Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	62%	> 90%	✘	10%	≥ 30%
Coverage of antenatal tetanus vaccination	50%	> 95%	✘	HIS	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	85%	≥ 90%	⚠	Total number of reported rapes	0
Proportion of births conducted by caesarean section	0%	5 - 15%	✘	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	4%	< 15%	✔	Proportion of eligible rape survivors provided with ECP within 120 hours	100%
Proportion of teenage pregnancies	1%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate ( / 1000 total births / month)	0			Number of genital ulcer diseases	4
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	2
Proportion of maternal deaths investigated within 48 hours		100%			

## HIV/AIDS

### PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.06	> 0.5
Proportion of TB patients tested for HIV		100%

### PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	12		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✗
PMTCT coverage	50%	100%	✗

## Nutrition

### NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	4%	< 20%
Prevalence of stunting (6-59 months)	24%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	0%	< 3%
Prevalence of anaemia in children (6-59 months)	29%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	13%	< 20%

### INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	59%	> 80%	✗
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	76%	> 70%	✗
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	

### MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/BBF)		> 70%

### CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	19	
Recovery rate	55%	> 75%
Default rate	38%	< 0%
Death rate	0%	< 10%
Referral rate	7%	

### MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	94		
Recovery rate	77%	> 75%	✓
Default rate	4%	< 15%	✓
Death rate	0%	< 3%	✓
Referral rate	0%		

## Water, Sanitation and Hygiene (WASH)

### WATER

	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	15	< 20
Proportion of households collecting >= 15 litres per person per day		> 95%
Number of persons of concern per water tap		< 80
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5		> 95%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%
Proportion of households collecting drinking water from protected water sources only		> 95%
Proportion of households with sufficient daily water storage capacity		> 80%

### WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters	20	1 : < 20,000	✓

### SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	73	< 20	✗
Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of households report defecating in a toilet		> 85%	
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	