

Site report 2018 Jewi ETHIOPIA

Population

42,770 refugees







Origin of refugees

SOUTH SUDAN

VARIOUS

Site opened Mar 2015

Partners











Key Indicators

CRUDE MORTALITY RATE

0.0

Deaths/1,000 refugees/

GAM

6% Global Acute **U5 MORTALITY RATE**

Deaths/1,000 refugees/

SAM

1%

Severe Acute Malnutrition

MATERNAL MORTALITY

Deaths

WATER SUPPLY

16

Liters/ person/ day


SKILLED BIRTH **ATTENDANCE**


of all births


LATRINE COVERAGE

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES








NON-COMMUNICABLE DISEASES





Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARI		
Number of medical doctors	1	1:<50,000	A	Incidence of malaria among children under 5	41			
Number of qualified nurses	16	1:<10,000	•	Incidence of watery diarrhoea among	11			
Number of community health workers	130	1:<1,000	•	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	29			
Consultations per trained clinician per day	31	< 50	•	Incidence of pneumonia among over 5	14			
Health utilisation rate (new visits / refugee /	0.4	1 - 4	8	Tuberculosis success rate	86%	< 90%	A	
year) Proportion of host population consultations	4%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	R STAN	IDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	②	Full vaccination coverage	75%	< !	95%	8
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5	②	Measles vaccination coverage	92%	< !	95%	8
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	②	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STAN	IDARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	1.1			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	9.9	50	- 150	8

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	58%	> 90%	8	Contraceptive prevalence rate	7%	≥ 30%	8
Coverage of antenatal tetanus vaccination	79%	> 95%	<u> </u>	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	99%	≥ 90%	②	Total number of reported rapes	8		
Proportion of births conducted by caesarean section	0%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours	100%	100%	•
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	1%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours	20%	100%	8
Proportion of teenage pregnancies	3%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	276		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	136		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	40	
Condom distribution rate	0.09	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	50%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	6%	< 20%	②	Proportion of children 0-23 months who were put to the breast within one hour of birth	91%	> 80%	②
Prevalence of stunting (6-59 months)	9%	< 20%	Ø	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	98%	> 70%	Ø
Prevalence of severe acute malnutrition (6 - 59 months)	1%	< 3%	②	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	51%	< 20%	8	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	37%	< 20%	8	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	100%	> 70%	Ø
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	647			Number of new moderate acute admissions (6-59 months)	1752		
Recovery rate	94%	> 75%	②	Recovery rate	92%	> 75%	②
Default rate	2%	< 0%	8	Default rate	3%	< 15%	②
Death rate	0%	< 10%	②	Death rate	0%	< 3%	②
Referral rate	1%			Referral rate	2%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	16	< 20	8	Number of hygiene promoters	60	1:<20,000	•
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	•	SANITATION Refugees per latrine/toilet	INDICATOR 11	STANDARD < 20	•
Number of persons of concern per water tap	140	< 80	8	Proportion of households with drop-hole latrine or drop-hole toilet	45%		
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%	•	Proportion of households report defecating in a toilet	100%	> 85%	•
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	8	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%	•
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	•				
Proportion of households with sufficient daily water storage capacity	85%	> 80%	②				