

# Site report 2018 Kebribeyah ETHIOPIA



## Population


**14,685** refugees

**19%**

Children under 5

**20%**

Women of reproductive age



Origin of refugees

**SOMALIA**

Site opened  
Jan 1991

## Partners



## Key Indicators

CRUDE MORTALITY RATE

**0.0** Deaths/1,000 refugees/month

U5 MORTALITY RATE

**0.1** Deaths/1,000 refugees/month

MATERNAL MORTALITY

**0** Deaths

SKILLED BIRTH ATTENDANCE

**100%** of all births

GAM

**NC** Global Acute Malnutrition

SAM

**NC** Severe Acute Malnutrition


WATER SUPPLY

**12** Liters/ person/ day



LATRINE COVERAGE

**12** Refugees/communal toilets or latrines


### COMMUNICABLE DISEASES




### INJURIES



### NON-COMMUNICABLE DISEASES



### MENTAL HEALTH



## Public Health

### HEALTH STAFFING

|                                    | INDICATOR | STANDARD     |
|------------------------------------|-----------|--------------|
| Number of medical doctors          |           | 1 : < 50,000 |
| Number of qualified nurses         |           | 1 : < 10,000 |
| Number of community health workers |           | 1 : < 1,000  |

### ACCESS AND UTILIZATION

|   | INDICATOR | STANDARD |
|---|-----------|----------|
| Consultations per trained clinician per day           | 42        | < 50     |
| Health utilisation rate (new visits / refugee / year) | 0.8       | 1 - 4    |
| Proportion of host population consultations           | 20%       |          |

### MORTALITY

|   | INDICATOR | STANDARD |
|---|-----------|----------|
| Crude Mortality Rate (CMR) (/1000/month)          | 0.0       | < 0.75   |
| Under-five Mortality Rate (U5MR) (/1000/month)    | 0.1       | < 1.5    |
| Infant Mortality Rate (IMR) (/1000 livebirths)    | 4.5       | < 30     |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths) | 0         | < 20     |

### OUTBREAK ALERT AND RESPONSE

|  | INDICATOR | STANDARD |
|--|-----------|----------|
| Proportion of outbreaks investigated within 48 hours |           | 100%     |
| Number of outbreaks reported                         | 0         |          |

### MORBIDITY

|  | INDICATOR | STANDARD |
|--|-----------|----------|
| Incidence of malaria among children under 5          | 0         |          |
| Incidence of watery diarrhoea among children under 5 | 24        |          |
| Incidence of pneumonia among children under 5        | 35        |          |
| Incidence of pneumonia among over 5                  | 13        |          |
| Tuberculosis success rate                            |           | < 90%    |
| Were any MDR/X-TB cases diagnosed among PoCs?        |           | No       |

### VACCINATION

|                              | INDICATOR | STANDARD |
|------------------------------|-----------|----------|
| Full vaccination coverage    | 68%       | < 95%    |
| Measles vaccination coverage | 68%       | < 95%    |
| Source of vaccination data   |           |          |

### IN-PATIENT DEPARTMENT (IPD)

|                               | INDICATOR | STANDARD |
|-------------------------------|-----------|----------|
| Average length of stay (days) | 5.3       |          |
| Case fatality rate            | 0.0       |          |
| Hospitalisation rate          | 2.0       | 50 - 150 |

## Reproductive Health

### ANTENATAL CARE

|   | INDICATOR | STANDARD |
|---|-----------|----------|
| Antenatal care coverage                   | 99%       | > 90%    |
| Coverage of antenatal tetanus vaccination | 99%       | > 95%    |

### DELIVERY CARE

|  | INDICATOR | STANDARD |
|--|-----------|----------|
| Proportion of births attended by skilled personnel                                     | 100%      | ≥ 90%    |
| Proportion of births conducted by caesarean section                                    | 1%        | 5 - 15%  |
| Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs) | 3%        | < 15%    |
| Proportion of teenage pregnancies  | 2%        |          |
| Still birth rate ( / 1000 total births / month)  | 1         |          |
| Number of maternal deaths  | 0         |          |
| Proportion of maternal deaths investigated within 48 hours                             |           | 100%     |

### FAMILY PLANNING

|   | INDICATOR | STANDARD |
|---|-----------|----------|
| Contraceptive prevalence rate           | 3%        | ≥ 30%    |
| Source of contraceptive prevalence data |           |          |

### SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

|  | INDICATOR | STANDARD |
|--|-----------|----------|
| Total number of reported rapes   | 0         |          |
| Proportion of eligible rape survivors provided with PEP within 72 hours  |           | 100%     |
| Proportion of eligible rape survivors provided with ECP within 120 hours |           | 100%     |

### SEXUALLY TRANSMITTED INFECTIONS (STIS)

|  | INDICATOR | STANDARD |
|--|-----------|----------|
| Number of genital ulcer diseases       | 7         |          |
| Number of Pelvic Inflammatory diseases | 15        |          |

## HIV/AIDS

### PREVENTION & TREATMENT

Proportion of blood units screened for HIV

INDICATOR STANDARD

100%

Condom distribution rate

0.02 > 0.5

Proportion of TB patients tested for HIV

100%

### PREVENTION & TREATMENT

Number of PoCs receiving ART

INDICATOR STANDARD

11

✘ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

95% 100%



## Nutrition

### NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

INDICATOR STANDARD

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

### INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

INDICATOR STANDARD

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

### MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

INDICATOR STANDARD

> 70%

### CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

INDICATOR STANDARD

27

Recovery rate

89% > 75%

Default rate

6% < 0%

Death rate

0% < 10%

Referral rate

6%

### MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

INDICATOR STANDARD

110

✔ Recovery rate

100% > 75%

✘ Default rate

0% < 15%

✔ Death rate

0% < 3%

Referral rate

0%

## Water, Sanitation and Hygiene (WASH)

### WATER

Average number of litres of potable water available per person per day

INDICATOR STANDARD

12 < 20

Proportion of households collecting >= 15 litres per person per day

> 95%

Number of persons of concern per water tap

< 80

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5

> 95%

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml

> 95%

Proportion of households collecting drinking water from protected water sources only

> 95%

Proportion of households with sufficient daily water storage capacity

> 80%

### WATER STAFFING

✘ Number of hygiene promoters

INDICATOR STANDARD

1 : < 20,000

### SANITATION

Refugees per latrine/toilet

INDICATOR STANDARD

12 < 20

Proportion of households with drop-hole latrine or drop-hole toilet

> 85%

Proportion of households report defecating in a toilet

> 90%

Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

> 90%