

Site report 2018 Kule ETHIOPIA



Population

53,008 refugees

19%

Children under 5

20%

Women of reproductive age



Origin of refugees

SOUTH SUDAN

Site opened

Feb 2015

Partners



Key Indicators

CRUDE MORTALITY RATE

0.2

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.3

Deaths/1,000 refugees/month

MATERNAL MORTALITY

0

Deaths

SKILLED BIRTH ATTENDANCE

98%

of all births

GAM

18%

Global Acute Malnutrition

SAM

3%

Severe Acute Malnutrition

WATER SUPPLY

16


Liters/ person/ day

LATRINE COVERAGE


10

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES





INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			INDICATOR	STANDARD	MORBIDITY			INDICATOR	STANDARD
Number of medical doctors	4	1 : < 50,000	✓	Incidence of malaria among children under 5	26				
Number of qualified nurses	96	1 : < 10,000	✓	Incidence of watery diarrhoea among children under 5	79				
Number of community health workers	200	1 : < 1,000	✓	Incidence of pneumonia among children under 5	85				
ACCESS AND UTILIZATION			INDICATOR	STANDARD	VACCINATION			INDICATOR	STANDARD
Consultations per trained clinician per day	32	< 50	✓	Incidence of pneumonia among over 5	34				
Health utilisation rate (new visits / refugee / year)	2.7	1 - 4	✓	Tuberculosis success rate	86%	< 90%	⚠		
Proportion of host population consultations	0%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	✗		
MORTALITY			INDICATOR	STANDARD	IN-PATIENT DEPARTMENT (IPD)			INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75	✓	Average length of stay (days)	8.4				
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5	✓	Case fatality rate	0.0				
Infant Mortality Rate (IMR) (/1000 livebirths)	9.9	< 30	✓	Hospitalisation rate	37.4	50 - 150	✗		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	✓						
OUTBREAK ALERT AND RESPONSE			INDICATOR	STANDARD					
Proportion of outbreaks investigated within 48 hours		100%							
Number of outbreaks reported	0								

Reproductive Health

ANTENATAL CARE			INDICATOR	STANDARD	FAMILY PLANNING			INDICATOR	STANDARD
Antenatal care coverage	21%	> 90%	✗	Contraceptive prevalence rate	9%	≥ 30%	✗		
Coverage of antenatal tetanus vaccination	97%	> 95%	✓	Source of contraceptive prevalence data	HIS				
DELIVERY CARE			INDICATOR	STANDARD	SEXUAL AND GENDER-BASED VIOLENCE (SGBV)			INDICATOR	STANDARD
Proportion of births attended by skilled personnel	98%	≥ 90%	✓	Total number of reported rapes	0				
Proportion of births conducted by caesarean section	0%	5 - 15%	✗	Proportion of eligible rape survivors provided with PEP within 72 hours	NR				
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	7%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours		100%			
Proportion of teenage pregnancies	8%			SEXUALLY TRANSMITTED INFECTIONS (STIS)			INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	866				
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	438				
Proportion of maternal deaths investigated within 48 hours		100%							

HIV/AIDS

PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.07	> 0.5
Proportion of TB patients tested for HIV		100%

PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	163		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✗
PMTCT coverage	102%	100%	✓

Nutrition

NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	18%	< 20%
Prevalence of stunting (6-59 months)	16%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	3%	< 3%
Prevalence of anaemia in children (6-59 months)	41%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	32%	< 20%

INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	89%	> 80%	✓
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	86%	> 70%	✓
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	✗

MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD	
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	100%	> 70%	✓

CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	1278	
Recovery rate	79%	> 75%
Default rate	4%	< 0%
Death rate	3%	< 10%
Referral rate	14%	

MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	2097		
Recovery rate	95%	> 75%	✓
Default rate	2%	< 15%	✓
Death rate	0%	< 3%	✓
Referral rate	2%		

Water, Sanitation and Hygiene (WASH)

WATER

	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	16	< 20
Proportion of households collecting >= 15 litres per person per day	100%	> 95%
Number of persons of concern per water tap	137	< 80
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%
Proportion of households collecting drinking water from protected water sources only	100%	> 95%
Proportion of households with sufficient daily water storage capacity	80%	> 80%

WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters	79	1 : < 20,000	✓

SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	10	< 20	✓
Proportion of households with drop-hole latrine or drop-hole toilet	46%		✗
Proportion of households report defecating in a toilet	95%	> 85%	✓
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%	✓