

Site report 2018 Melkadida ETHIOPIA



Population


34,223 refugees

13%

Children under 5

20%

Women of reproductive age



Origin of refugees

SOMALIA

Site opened
Feb 2010

Partners



Key Indicators

CRUDE MORTALITY RATE

0.2 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.6 Deaths/1,000 refugees/month

MATERNAL MORTALITY

3 Deaths

SKILLED BIRTH ATTENDANCE

106% of all births

GAM

10% Global Acute Malnutrition

SAM

2% Severe Acute Malnutrition


WATER SUPPLY

20 Liters/ person/ day



LATRINE COVERAGE

29 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000
Number of qualified nurses		1 : < 10,000
Number of community health workers		1 : < 1,000

ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	45	< 50
Health utilisation rate (new visits / refugee / year)	1.2	1 - 4
Proportion of host population consultations	19%	

MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.6	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	17.6	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	11	< 20

OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%
Number of outbreaks reported	0	

MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	0	
Incidence of watery diarrhoea among children under 5	30	
Incidence of pneumonia among children under 5	13	
Incidence of pneumonia among over 5	4	
Tuberculosis success rate		< 90%
Were any MDR/X-TB cases diagnosed among PoCs?		No

VACCINATION

	INDICATOR	STANDARD
Full vaccination coverage	39%	< 95%
Measles vaccination coverage	100%	< 95%
Source of vaccination data		

IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD
Average length of stay (days)	4.6	
Case fatality rate	0.0	
Hospitalisation rate	7.1	50 - 150

Reproductive Health

ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	33%	> 90%
Coverage of antenatal tetanus vaccination	59%	> 95%

DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	106%	≥ 90%
Proportion of births conducted by caesarean section	0%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	4%	< 15%
Proportion of teenage pregnancies	3%	
Still birth rate (/ 1000 total births / month)	1	
Number of maternal deaths	3	
Proportion of maternal deaths investigated within 48 hours	100%	100%

FAMILY PLANNING

	INDICATOR	STANDARD
Contraceptive prevalence rate	1%	≥ 30%
Source of contraceptive prevalence data		

SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD
Total number of reported rapes	0	
Proportion of eligible rape survivors provided with PEP within 72 hours		100%
Proportion of eligible rape survivors provided with ECP within 120 hours		100%

SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD
Number of genital ulcer diseases	9	
Number of Pelvic Inflammatory diseases	5	

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receiving ART	7	
Condom distribution rate	0.07	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	89%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	10%	< 20%	Proportion of children 0-23 months who were put to the breast within one hour of birth	85%	> 80%
Prevalence of stunting (6-59 months)	25%	< 20%	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	73%	> 70%
Prevalence of severe acute malnutrition (6 - 59 months)	2%	< 3%	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	45%	< 20%			
Prevalence of anaemia in women of reproductive age (15-49 yrs)	37%	< 20%			

CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	354		Number of new moderate acute admissions (6-59 months)	909	
Recovery rate	96%	> 75%	Recovery rate	95%	> 75%
Default rate	2%	< 0%	Default rate	2%	< 15%
Death rate	0%	< 10%	Death rate	0%	< 3%
Referral rate	2%		Referral rate	3%	

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	20	< 20	Number of hygiene promoters		1 : < 20,000
Proportion of households collecting >= 15 litres per person per day		> 95%			
Number of persons of concern per water tap		< 80			
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%			
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%			
Proportion of households collecting drinking water from protected water sources only		> 95%			
Proportion of households with sufficient daily water storage capacity		> 80%			

SANITATION	INDICATOR	STANDARD
Refugees per latrine/toilet	29	< 20
Proportion of households with drop-hole latrine or drop-hole toilet		
Proportion of households report defecating in a toilet		> 85%
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%