



Site report 2018 Nguenyil ETHIOPIA



Population

74,596 refugees



Origin of refugees
SOUTH SUDAN

Site opened
Oct 2016

Partners



Key Indicators

CRUDE MORTALITY RATE

0.0 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.0 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

82% of all births

GAM

14% Global Acute Malnutrition

SAM

3% Severe Acute Malnutrition


WATER SUPPLY

13 Liters/ person/ day


LATRINE COVERAGE

13 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES





INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	1	1 : < 50,000	✘	12	
Number of qualified nurses	12	1 : < 10,000	✔	26	
Number of community health workers	200	1 : < 1,000	✔	21	
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	47	< 50	✔	19%	< 95%
Health utilisation rate (new visits / refugee / year)	1.2	1 - 4	✔	90%	< 95%
Proportion of host population consultations	0%		✔	HIS	
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	✔	Average length of stay (days)	4.0
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5	✔	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	✔	Hospitalisation rate	6.9
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	✔		50 - 150
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%			
Number of outbreaks reported	0				

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	32%	> 90%	✘	1%	≥ 30%
Coverage of antenatal tetanus vaccination	63%	> 95%	⚠	HIS	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	82%	≥ 90%	⚠	Total number of reported rapes	3
Proportion of births conducted by caesarean section	0%	5 - 15%	✘	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	1%	< 15%	✔	Proportion of eligible rape survivors provided with ECP within 120 hours	100%
Proportion of teenage pregnancies	3%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	0
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	3
Proportion of maternal deaths investigated within 48 hours		100%			

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receiving ART	0	
Condom distribution rate	0.14	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	100%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	14%	< 20%	Proportion of children 0-23 months who were put to the breast within one hour of birth	69%	> 80%
Prevalence of stunting (6-59 months)	10%	< 20%	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	66%	> 70%
Prevalence of severe acute malnutrition (6 - 59 months)	3%	< 3%	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	38%	< 20%			
Prevalence of anaemia in women of reproductive age (15-49 yrs)	29%	< 20%			

CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	769		Number of new moderate acute admissions (6-59 months)	1581	
Recovery rate	89%	> 75%	Recovery rate	89%	> 75%
Default rate	4%	< 0%	Default rate	1%	< 15%
Death rate	0%	< 10%	Death rate	0%	< 3%
Referral rate	4%		Referral rate	8%	

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	13	< 20	Number of hygiene promoters	170	1 : < 20,000
Proportion of households collecting >= 15 litres per person per day	40%	> 95%			
Number of persons of concern per water tap	133	< 80			
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%			
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%			
Proportion of households collecting drinking water from protected water sources only	100%	> 95%			
Proportion of households with sufficient daily water storage capacity	95%	> 80%			

SANITATION	INDICATOR	STANDARD
Refugees per latrine/toilet	13	< 20
Proportion of households with drop-hole latrine or drop-hole toilet	35%	
Proportion of households report defecating in a toilet	90%	> 85%
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%