

Site report 2018 Okugo ETHIOPIA



Population


14,082 refugees

28%

Children under 5

20%

Women of reproductive age



Origin of refugees

SOUTH SUDAN

Site opened

Apr 2014

Partners



Key Indicators

CRUDE MORTALITY RATE

0.0

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.0

Deaths/1,000 refugees/month

MATERNAL MORTALITY

0

Deaths

SKILLED BIRTH ATTENDANCE

87%

of all births

GAM

9%

Global Acute Malnutrition

SAM

1%

Severe Acute Malnutrition

WATER SUPPLY

18


Liters/ person/ day

LATRINE COVERAGE


11

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES





INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	1	1 : < 50,000	✓	5	
Number of qualified nurses	7	1 : < 10,000	✓	23	
Number of community health workers	36	1 : < 1,000	✓	18	
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	23	< 50	✓	31%	< 95%
Health utilisation rate (new visits / refugee / year)	1.1	1 - 4	✓	84%	< 95%
Proportion of host population consultations	14%		✓	HIS	
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	✓	Average length of stay (days)	2.6
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5	✓	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	3.5	< 30	✓	Hospitalisation rate	10.4
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	✓		50 - 150
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%			
Number of outbreaks reported	0				

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	71%	> 90%	✗	7%	≥ 30%
Coverage of antenatal tetanus vaccination	74%	> 95%	⚠	Source of contraceptive prevalence data	HIS
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	87%	≥ 90%	⚠	Total number of reported rapes	0
Proportion of births conducted by caesarean section	0%	5 - 15%	✗	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	4%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%
Proportion of teenage pregnancies	7%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	5
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	2
Proportion of maternal deaths investigated within 48 hours		100%			

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receiving ART	153	
Condom distribution rate	0.37	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	89%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	9%	< 20%	Proportion of children 0-23 months who were put to the breast within one hour of birth	72%	> 80%
Prevalence of stunting (6-59 months)	5%	< 20%	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	65%	> 70%
Prevalence of severe acute malnutrition (6 - 59 months)	1%	< 3%	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	43%	< 20%	MICRONUTRIENT DEFICIENCY		
Prevalence of anaemia in women of reproductive age (15-49 yrs)	20%	< 20%	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	100%	> 70%

CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	69		Number of new moderate acute admissions (6-59 months)	129	
Recovery rate	91%	> 75%	Recovery rate	96%	> 75%
Default rate	2%	< 0%	Default rate	1%	< 15%
Death rate	2%	< 10%	Death rate	0%	< 3%
Referral rate	5%		Referral rate	0%	

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	18	< 20	Number of hygiene promoters	11	1 : < 20,000
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	SANITATION		
Number of persons of concern per water tap	111	< 80	Refugees per latrine/toilet	11	< 20
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%	Proportion of households with drop-hole latrine or drop-hole toilet	31%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	Proportion of households report defecating in a toilet	90%	> 85%
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	90%	> 90%
Proportion of households with sufficient daily water storage capacity	80%	> 80%			