



Site report 2018 Sherkole ETHIOPIA



Population

8,968 refugees



Origin of refugees

SUDAN CONGO BURUNDI
RWANDA

Site opened
Jan 1997

Partners



Key Indicators

CRUDE MORTALITY RATE

0.2 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.5 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

100% of all births

GAM

9% Global Acute Malnutrition

SAM

1% Severe Acute Malnutrition


WATER SUPPLY

22 Liters/ person/ day



LATRINE COVERAGE

6 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors	1	1 : < 50,000
Number of qualified nurses	12	1 : < 10,000
Number of community health workers	62	1 : < 1,000

ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	38	< 50
Health utilisation rate (new visits / refugee / year)	1.7	1 - 4
Proportion of host population consultations	10%	

MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.5	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	33.0	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	14	< 20

OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%
Number of outbreaks reported	0	

MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	1	
Incidence of watery diarrhoea among children under 5	43	
Incidence of pneumonia among children under 5	89	
Incidence of pneumonia among over 5	28	
Tuberculosis success rate		< 90%
Were any MDR/X-TB cases diagnosed among PoCs?	No	No

VACCINATION

	INDICATOR	STANDARD
Full vaccination coverage	62%	< 95%
Measles vaccination coverage	93%	< 95%
Source of vaccination data	HIS	

IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD
Average length of stay (days)	5.6	
Case fatality rate	0.0	
Hospitalisation rate	12.9	50 - 150

Reproductive Health

ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	94%	> 90%
Coverage of antenatal tetanus vaccination	95%	> 95%

DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	100%	≥ 90%
Proportion of births conducted by caesarean section	0%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	2%	< 15%
Proportion of teenage pregnancies	2%	
Still birth rate (/ 1000 total births / month)	4	
Number of maternal deaths	0	
Proportion of maternal deaths investigated within 48 hours		100%

FAMILY PLANNING

	INDICATOR	STANDARD
Contraceptive prevalence rate	14%	≥ 30%
Source of contraceptive prevalence data	HIS	

SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD
Total number of reported rapes	2	
Proportion of eligible rape survivors provided with PEP within 72 hours		100%
Proportion of eligible rape survivors provided with ECP within 120 hours		100%

SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD
Number of genital ulcer diseases	1	
Number of Pelvic Inflammatory diseases	0	

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receiving ART	52	
Condom distribution rate	0.57	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	102%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	9%	< 20%	Proportion of children 0-23 months who were put to the breast within one hour of birth	87%	> 80%
Prevalence of stunting (6-59 months)	26%	< 20%	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	88%	> 70%
Prevalence of severe acute malnutrition (6 - 59 months)	1%	< 3%	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	20%	< 20%			
Prevalence of anaemia in women of reproductive age (15-49 yrs)	9%	< 20%			
CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	0		Number of new moderate acute admissions (6-59 months)	20	
Recovery rate		> 75%	Recovery rate	96%	> 75%
Default rate		< 0%	Default rate	0%	< 15%
Death rate		< 10%	Death rate	4%	< 3%
Referral rate			Referral rate	0%	

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	22	< 20	Number of hygiene promoters	18	1 : < 20,000
Proportion of households collecting >= 15 litres per person per day	100%	> 95%			
Number of persons of concern per water tap	75	< 80			
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%			
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%			
Proportion of households collecting drinking water from protected water sources only	100%	> 95%			
Proportion of households with sufficient daily water storage capacity		> 80%			

SANITATION	INDICATOR	STANDARD
Refugees per latrine/toilet	6	< 20
Proportion of households with drop-hole latrine or drop-hole toilet	8%	
Proportion of households report defecating in a toilet	47%	> 85%
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	11%	> 90%