



# Site report 2018 Shimelba ETHIOPIA



## Population

**7,732** refugees



Origin of refugees

**ERITREA**

Site opened

**Jan 2004**

## Partners



## Key Indicators

CRUDE MORTALITY RATE

**0** Deaths/1,000 refugees/month

U5 MORTALITY RATE

**0** Deaths/1,000 refugees/month

MATERNAL MORTALITY

**0** Deaths

SKILLED BIRTH ATTENDANCE

**100%** of all births

GAM

**15%** Global Acute Malnutrition

SAM

**3%** Severe Acute Malnutrition


WATER SUPPLY

**27** Liters/ person/ day

LATRINE COVERAGE


**7** Refugees/communal toilets or latrines

### COMMUNICABLE DISEASES




### INJURIES

### NON-COMMUNICABLE DISEASES



### MENTAL HEALTH



## Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	1	1 : < 50,000	✓	2	
Number of qualified nurses	9	1 : < 10,000	✓	19	
Number of community health workers	33	1 : < 1,000	✓	45	
ACCESS AND UTILIZATION			MORTALITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	28	< 50	✓	8	
Health utilisation rate (new visits / refugee / year)	2.1	1 - 4	✓	90%	< 90% ✓
Proportion of host population consultations	44%			No	No ✗
			✓	Incidence of malaria among children under 5	
			✓	Incidence of watery diarrhoea among children under 5	
			✓	Incidence of pneumonia among children under 5	
			✓	Incidence of pneumonia among over 5	
			✓	Tuberculosis success rate	
			✓	Were any MDR/X-TB cases diagnosed among PoCs?	
MORTALITY			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	✓	67%	< 95% ✗
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5	✓	67%	< 95% ✗
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	✓	SOURCE OF VACCINATION DATA	HIS
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	✓	IN-PATIENT DEPARTMENT (IPD)	
OUTBREAK ALERT AND RESPONSE				INDICATOR	STANDARD
	INDICATOR	STANDARD			
Proportion of outbreaks investigated within 48 hours		100%		5.5	
Number of outbreaks reported	0			0.0	
				8.1	50 - 150 ✗

## Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	100%	> 90%	✓	68%	≥ 30% ✓
Coverage of antenatal tetanus vaccination	100%	> 95%	✓	SOURCE OF CONTRACEPTIVE PREVALENCE DATA	HIS
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	100%	≥ 90%	✓	0	
Proportion of births conducted by caesarean section	5%	5 - 15%	✓	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	0%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%
Proportion of teenage pregnancies	1%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate ( / 1000 total births / month)	0			Number of genital ulcer diseases	0
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0
Proportion of maternal deaths investigated within 48 hours		100%			

## HIV/AIDS

### PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.55	> 0.5
Proportion of TB patients tested for HIV		100%

### PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	12		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✓
PMTCT coverage	115%	100%	✓

## Nutrition

### NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	15%	< 20%
Prevalence of stunting (6-59 months)	23%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	3%	< 3%
Prevalence of anaemia in children (6-59 months)	25%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	10%	< 20%

### INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	77%	> 80%	✗
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	68%	> 70%	✗
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	

### MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/BBF)		> 70%

### CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	13	
Recovery rate	100%	> 75%
Default rate	0%	< 0%
Death rate	0%	< 10%
Referral rate	0%	

### MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	45		
Recovery rate	92%	> 75%	✓
Default rate	0%	< 15%	✓
Death rate	0%	< 3%	✓
Referral rate	0%		

## Water, Sanitation and Hygiene (WASH)

### WATER

	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	27	< 20	✓
Proportion of households collecting >= 15 litres per person per day		> 95%	
Number of persons of concern per water tap		< 80	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5		> 95%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	
Proportion of households collecting drinking water from protected water sources only		> 95%	
Proportion of households with sufficient daily water storage capacity		> 80%	

### WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters	16	1 : < 20,000	✓

### SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	7	< 20	✓
Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of households report defecating in a toilet		> 85%	
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	