



Site report 2018 Tierkidi ETHIOPIA



Population

71,860 refugees



Origin of refugees
SOUTH SUDAN

Site opened
Feb 2014

Partners



Key Indicators

CRUDE MORTALITY RATE

0.0 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.4 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

99% of all births

GAM

11% Global Acute Malnutrition

SAM

2% Severe Acute Malnutrition


WATER SUPPLY

16 Liters/ person/ day



LATRINE COVERAGE

19 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors	1	1 : < 50,000
Number of qualified nurses	15	1 : < 10,000
Number of community health workers	140	1 : < 1,000

ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	44	< 50
Health utilisation rate (new visits / refugee / year)	1.3	1 - 4
Proportion of host population consultations	1%	

MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.4	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	3.9	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20

OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%
Number of outbreaks reported	0	

MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	68	
Incidence of watery diarrhoea among children under 5	121	
Incidence of pneumonia among children under 5	87	
Incidence of pneumonia among over 5	13	
Tuberculosis success rate	88%	< 90% ⚠️
Were any MDR/X-TB cases diagnosed among PoCs?	No	No ❌

VACCINATION

	INDICATOR	STANDARD
Full vaccination coverage	35%	< 95% ❌
Measles vaccination coverage	94%	< 95% ❌
Source of vaccination data	HIS	

IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD
Average length of stay (days)	5.3	
Case fatality rate	0.0	
Hospitalisation rate	8.0	50 - 150 ❌

Reproductive Health

ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	65%	> 90%
Coverage of antenatal tetanus vaccination	77%	> 95%

DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	99%	≥ 90%
Proportion of births conducted by caesarean section	0%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	2%	< 15%
Proportion of teenage pregnancies	2%	
Still birth rate (/ 1000 total births / month)	0	
Number of maternal deaths	0	
Proportion of maternal deaths investigated within 48 hours		100%

FAMILY PLANNING

	INDICATOR	STANDARD
Contraceptive prevalence rate	15%	≥ 30% ❌
Source of contraceptive prevalence data	HIS	

SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD
Total number of reported rapes	1	
Proportion of eligible rape survivors provided with PEP within 72 hours	0%	100% ❌
Proportion of eligible rape survivors provided with ECP within 120 hours	0%	100% ❌

SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD
Number of genital ulcer diseases	160	
Number of Pelvic Inflammatory diseases	49	

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receiving ART	123	
Condom distribution rate	0.04	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	93%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	11%	< 20%	Proportion of children 0-23 months who were put to the breast within one hour of birth	94%	> 80%
Prevalence of stunting (6-59 months)	15%	< 20%	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	92%	> 70%
Prevalence of severe acute malnutrition (6 - 59 months)	2%	< 3%	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	39%	< 20%	MICRONUTRIENT DEFICIENCY		
Prevalence of anaemia in women of reproductive age (15-49 yrs)	35%	< 20%	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	100%	> 70%

CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	515		Number of new moderate acute admissions (6-59 months)	1084	
Recovery rate	93%	> 75%	Recovery rate	94%	> 75%
Default rate	5%	< 0%	Default rate	4%	< 15%
Death rate	0%	< 10%	Death rate	0%	< 3%
Referral rate	1%		Referral rate	1%	

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	16	< 20	Number of hygiene promoters	55	1 : < 20,000
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	SANITATION		
Number of persons of concern per water tap	232	< 80	Refugees per latrine/toilet	19	< 20
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%	Proportion of households with drop-hole latrine or drop-hole toilet	23%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	Proportion of households report defecating in a toilet	95%	> 85%
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%
Proportion of households with sufficient daily water storage capacity	80%	> 80%			