

Site report 2018 Tongo ETHIOPIA

Population

12,572 refugees





The UN Refugee Agency

Origin of refugees

SOUTH SUDAN SUDAN

Site opened Nov 2011

Partners







Key Indicators

CRUDE MORTALITY RATE

0.0

Deaths/1,000 refugees/ month

GAM

Global Acute

U5 MORTALITY RATE

Deaths/1,000 refugees/ month

SAM

2%

Severe Acute

MATERNAL MORTALITY

Deaths

WATER SUPPLY

19

Liters/ person/ day


SKILLED BIRTH ATTENDANCE

100% of all births


LATRINE COVERAGE


Refugees/communal

COMMUNICABLE DISEASES




INJURIES






NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STAND	ARD	
Number of medical doctors	1	1:<50,000	•	Incidence of malaria among children under 5	0			
Number of qualified nurses	10	1:<10,000	•	Incidence of watery diarrhoea among	30			
Number of community health workers	56	1:<1,000	②	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	23			
Consultations per trained clinician per day	49	< 50	•	Incidence of pneumonia among over 5	8			
Health utilisation rate (new visits / refugee /	1.2	1 - 4	•	Tuberculosis success rate		< 90	%	
year) Proportion of host population consultations	35%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	R S	TANDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	②	Full vaccination coverage	83%		< 95%	8
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5	②	Measles vaccination coverage	92%		< 95%	8
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	②	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATO	R S	TANDARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	6.4			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	14.2		50 - 150	8

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	88%	> 90%	<u> </u>	Contraceptive prevalence rate	33%	≥ 30%	②
Coverage of antenatal tetanus vaccination	95%	> 95%	<u> </u>	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	100%	≥ 90%	②	Total number of reported rapes	2		
Proportion of births conducted by caesarean section	0%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours	0%	100%	8
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	1%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours	0%	100%	8
Proportion of teenage pregnancies	1%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	0		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	10	
Condom distribution rate	0.36	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	96%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	13%	< 20%	A	Proportion of children 0-23 months who were put to the breast within one hour of birth	83%	> 80%	②
Prevalence of stunting (6-59 months)	42%	< 20%	②	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	88%	> 70%	②
Prevalence of severe acute malnutrition (6 - 59 months)	2%	< 3%	②	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	30%	< 20%	8	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	11%	< 20%	Ø	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	100%	> 70%	Ø
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	38			Number of new moderate acute admissions (6-59 months)	148		
Recovery rate	100%	> 75%	②	Recovery rate	98%	> 75%	②
Default rate	0%	< 0%	8	Default rate	0%	< 15%	②
Death rate	0%	< 10%	②	Death rate	1%	< 3%	②
Referral rate	0%			Referral rate	1%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	19	< 20	A	Number of hygiene promoters	15	1:<20,000	•
Proportion of households collecting >= 15	100%	> 95%	•	SANITATION	INDICATOR	STANDARD	
litres per person per day				Refugees per latrine/toilet	9	< 20	Ø
Number of persons of concern per water tap	0	< 80	②	Proportion of households with drop-hole latrine or drop-hole toilet	53%		
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%	②	Proportion of households report defecating in a toilet	73%	> 85%	A
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	8	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	17%	> 90%	8
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	•				
Proportion of households with sufficient daily water storage capacity		> 80%					