

Site report 2018 Azraq JORDAN



Population


36,389 refugees

22%

Children under 5

23%

Women of reproductive age



Origin of refugees

SYRIA

Site opened
May 2014

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.3

Deaths/1,000 refugees/month

MATERNAL MORTALITY

0

Deaths

SKILLED BIRTH ATTENDANCE

97%

of all births

GAM

NC

Global Acute Malnutrition

SAM

NC

Severe Acute Malnutrition

WATER SUPPLY

35

Liters/ person/ day

LATRINE COVERAGE

15

Refugees/communal toilets or latrines

Public Health

HEALTH STAFFING	INDICATOR	STANDARD	MORTALITY	INDICATOR	STANDARD	
Number of medical doctors	41	1 : < 50,000	✓	Crude Mortality Rate (CMR) (/1000/month)	0.1 < 0.75 ✓	
Number of qualified nurses	48	1 : < 10,000	✓	Under-five Mortality Rate (U5MR) (/1000/month)	0.3 < 1.5 ✓	
Number of community health workers	38	1 : < 1,000	✓	Infant Mortality Rate (IMR) (/1000 livebirths)	19.3 < 30 ✓	
ACCESS AND UTILIZATION	INDICATOR	STANDARD	Neonatal Mortality Rate (NNMR) (/1000 livebirths)	14	< 20 ✓	
Consultations per trained clinician per day	46	< 50	✓	IN-PATIENT DEPARTMENT (IPD)	INDICATOR	STANDARD
Health utilisation rate (new visits / refugee / year)	8.2	1 - 4	✓	Average length of stay (days)	2.0	
Proportion of host population consultations	0%			Case fatality rate		
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD	✓	Hospitalisation rate	109.0	50 - 150 ✓
Proportion of outbreaks investigated within 48 hours	100%	100%				
Number of outbreaks reported	1					

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD	FAMILY PLANNING	INDICATOR	STANDARD
Antenatal care coverage	68%	> 90%	✗	Contraceptive prevalence rate	≥ 30%
Coverage of antenatal tetanus vaccination	38%	> 95%	✗	Source of contraceptive prevalence data	HIS
DELIVERY CARE	INDICATOR	STANDARD	SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	97%	≥ 90%	✓	Total number of reported rapes	4
Proportion of births conducted by caesarean section	19%	5 - 15%	✓	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	7%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%
Still birth rate (/ 1000 total births / month)	3				
Number of maternal deaths	0				
Proportion of maternal deaths investigated within 48 hours		100%			

Nutrition

NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

INDICATOR STANDARD

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

> 70%

CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

INDICATOR STANDARD

24

Recovery rate

93% > 75%



Default rate

3% < 0%



Death rate

3% < 10%



Referral rate

MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

INDICATOR STANDARD

153

Recovery rate

98% > 75%



Default rate

2% < 15%



Death rate

2% < 3%



Referral rate

Water, Sanitation and Hygiene (WASH)

WATER

Average number of litres of potable water available per person per day

INDICATOR STANDARD

35

< 20



Proportion of households collecting >= 15 litres per person per day

> 95%

Number of persons of concern per water tap

< 80

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

> 95%

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/100 ml

> 95%

Proportion of households collecting drinking water from protected water sources only

0% > 95%



Proportion of households with sufficient daily water storage capacity

0% > 80%



WATER STAFFING

Number of hygiene promoters

INDICATOR STANDARD

1 : < 20,000



SANITATION

Refugees per latrine/toilet

INDICATOR STANDARD

15

< 20



Proportion of households with drop-hole latrine or drop-hole toilet

Proportion of households report defecating in a toilet

0% > 85%



Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

0% > 90%

