

Site report 2018 Dagahaley KENYA



Population


69,552 refugees

17%

Children under 5

20%

Women of reproductive age



Origin of refugees

SOMALIA ETHIOPIA SOUTHERN SUDAN ERITREA

Site opened Jan 1992

Partners



Key Indicators

CRUDE MORTALITY RATE

0.0 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.2 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

81% of all births

GAM

9% Global Acute Malnutrition

SAM

2% Severe Acute Malnutrition


WATER SUPPLY

28 Liters/ person/ day



LATRINE COVERAGE

6 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	4	1 : < 50,000	✓	Incidence of malaria among children under 5	0
Number of qualified nurses	52	1 : < 10,000	✓	Incidence of watery diarrhoea among children under 5	24
Number of community health workers	152	1 : < 1,000	✓	Incidence of pneumonia among children under 5	32
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	37	< 50	✓	Incidence of pneumonia among over 5	13
Health utilisation rate (new visits / refugee / year)	1.0	1 - 4	⚠	Tuberculosis success rate	90% < 90% ⚠
Proportion of host population consultations	0%			Were any MDR/X-TB cases diagnosed among PoCs?	Yes No ✗
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	✓	Average length of stay (days)	2.2
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5	✓	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	15.5	< 30	✓	Hospitalisation rate	75.6 50 - 150 ✓
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	13	< 20	✓		
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%			
Number of outbreaks reported	0				

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	83%	> 90%	⚠	Contraceptive prevalence rate	5% ≥ 30% ✗
Coverage of antenatal tetanus vaccination	85%	> 95%	⚠	Source of contraceptive prevalence data	HIS
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	81%	≥ 90%	⚠	Total number of reported rapes	21
Proportion of births conducted by caesarean section	4%	5 - 15%	✗	Proportion of eligible rape survivors provided with PEP within 72 hours	100% 100% ✓
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	5%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	86% 100% ✗
Proportion of teenage pregnancies	4%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	3			Number of genital ulcer diseases	0
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	1
Proportion of maternal deaths investigated within 48 hours		100%			

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	✓ Number of PoCs receiving ART	36	
Condom distribution rate	0.00	> 0.5	✗ Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	100%	100% ✓

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	9%	< 20%	✓ Proportion of children 0-23 months who were put to the breast within one hour of birth	79%	> 80% ⚠
Prevalence of stunting (6-59 months)	27%	< 20%	✓ Proportion of infants aged between 0-5 months who received only breast milk during the previous day	63%	> 70% ⚠
Prevalence of severe acute malnutrition (6 - 59 months)	2%	< 3%	✓ IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	61%	< 20%	✗ MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD
Prevalence of anaemia in women of reproductive age (15-49 yrs)	53%	< 20%	✗ Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	61%	> 70% ⚠

CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	1205		Number of new moderate acute admissions (6-59 months)	2053	
Recovery rate	88%	> 75%	✓ Recovery rate	89%	> 75% ✓
Default rate	4%	< 0%	✗ Default rate	1%	< 15% ✓
Death rate	1%	< 10%	✓ Death rate	0%	< 3% ✓
Referral rate	2%		Referral rate	4%	

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	28	< 20	✓ Number of hygiene promoters	50	1 : < 20,000 ✓
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	✓	INDICATOR	STANDARD
Number of persons of concern per water tap	93	< 80	⚠	Refugees per latrine/toilet	6 < 20 ✓
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%	✓	Proportion of households with drop-hole latrine or drop-hole toilet	75%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	✗	Proportion of households report defecating in a toilet	99% > 85% ✓
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	✓	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	0% > 90% ✗
Proportion of households with sufficient daily water storage capacity	100%	> 80%	✓		