



Site report 2018 Hagadera KENYA



Population

74,050 refugees



Origin of refugees

SOMALIA ETHIOPIA NONE

Site opened
Jan 1992

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.2 Deaths/1,000 refugees/month

MATERNAL MORTALITY

5 Deaths

SKILLED BIRTH ATTENDANCE

94% of all births

GAM

6% Global Acute Malnutrition

SAM

1% Severe Acute Malnutrition


WATER SUPPLY

31 Liters/ person/ day



LATRINE COVERAGE

5 Refugees/communal toilets or latrines

COMMUNICABLE DISEASES




INJURIES




Total Consultations
79,235

NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors	5	1 : < 50,000
Number of qualified nurses	29	1 : < 10,000
Number of community health workers	110	1 : < 1,000

ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	93	< 50
Health utilisation rate (new visits / refugee / year)	0.9	1 - 4
Proportion of host population consultations	0%	

MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	12.5	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	11	< 20

OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours	100%	100%
Number of outbreaks reported	2	

MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	0	
Incidence of watery diarrhoea among children under 5	31	
Incidence of pneumonia among children under 5	23	
Incidence of pneumonia among over 5	10	
Tuberculosis success rate	95%	< 90%
Were any MDR/X-TB cases diagnosed among PoCs?	No	No

VACCINATION

	INDICATOR	STANDARD
Full vaccination coverage	52%	< 95%
Measles vaccination coverage	100%	< 95%
Source of vaccination data	survey	

IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD
Average length of stay (days)	3.5	
Case fatality rate	0.0	
Hospitalisation rate	57.4	50 - 150

Reproductive Health

ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	98%	> 90%
Coverage of antenatal tetanus vaccination	98%	> 95%

DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	94%	≥ 90%
Proportion of births conducted by caesarean section	6%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	6%	< 15%
Proportion of teenage pregnancies	3%	
Still birth rate (/ 1000 total births / month)	1	
Number of maternal deaths	5	
Proportion of maternal deaths investigated within 48 hours	100%	100%

FAMILY PLANNING

	INDICATOR	STANDARD
Contraceptive prevalence rate	8%	≥ 30%
Source of contraceptive prevalence data	HIS	

SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD
Total number of reported rapes	15	
Proportion of eligible rape survivors provided with PEP within 72 hours	91%	100%
Proportion of eligible rape survivors provided with ECP within 120 hours	83%	100%

SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD
Number of genital ulcer diseases	19	
Number of Pelvic Inflammatory diseases	88	

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	✓ Number of PoCs receiving ART	24	
Condom distribution rate	0.01	> 0.5	✗ Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	100%	100% ✓

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	6%	< 20%	✓ Proportion of children 0-23 months who were put to the breast within one hour of birth	85%	> 80% ✓
Prevalence of stunting (6-59 months)	19%	< 20%	✓ Proportion of infants aged between 0-5 months who received only breast milk during the previous day	67%	> 70% ⚠
Prevalence of severe acute malnutrition (6 - 59 months)	1%	< 3%	✓ IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	54%	< 20%	✗ MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD
Prevalence of anaemia in women of reproductive age (15-49 yrs)	48%	< 20%	✗ Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	61%	> 70% ⚠

CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	1122		Number of new moderate acute admissions (6-59 months)	4092	
Recovery rate	97%	> 75%	✓ Recovery rate	99%	> 75% ✓
Default rate	0%	< 0%	✗ Default rate	0%	< 15% ✓
Death rate	1%	< 10%	✓ Death rate	0%	< 3% ✓
Referral rate	1%		Referral rate	1%	

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	31	< 20	✓ Number of hygiene promoters	50	1 : < 20,000 ✓
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	✓ SANITATION	INDICATOR	STANDARD
Number of persons of concern per water tap	93	< 80	⚠ Refugees per latrine/toilet	5	< 20 ✓
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%	✓ Proportion of households with drop-hole latrine or drop-hole toilet	86%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	✓ Proportion of households report defecating in a toilet	98%	> 85% ✓
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	✗ Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90% ✓
Proportion of households with sufficient daily water storage capacity	100%	> 80%	✓		