



# Site report 2018 Ifo KENYA



## Population

**65,143** refugees



### Origin of refugees

SOMALIA ETHIOPIA SOUTHERN SUDAN DRC

Site opened Jan 1992

## Partners



## Key Indicators

### CRUDE MORTALITY RATE

**0.1** Deaths/1,000 refugees/month

### U5 MORTALITY RATE

**0.2** Deaths/1,000 refugees/month

### MATERNAL MORTALITY

**2** Deaths

### SKILLED BIRTH ATTENDANCE

**84%** of all births

### GAM

**NC** Global Acute Malnutrition

### SAM

**NC** Severe Acute Malnutrition


### WATER SUPPLY

Liters/ person/ day



### LATRINE COVERAGE

Refugees/communal toilets or latrines


## COMMUNICABLE DISEASES




## INJURIES



## NON-COMMUNICABLE DISEASES



## MENTAL HEALTH



## Public Health

### HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors	2	1 : < 50,000
Number of qualified nurses	24	1 : < 10,000
Number of community health workers	102	1 : < 1,000

### ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	73	< 50
Health utilisation rate (new visits / refugee / year)	1.1	1 - 4
Proportion of host population consultations	0%	

### MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	26.9	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	12	< 20

### OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%
Number of outbreaks reported	0	

### MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	0	
Incidence of watery diarrhoea among children under 5	41	
Incidence of pneumonia among children under 5	40	
Incidence of pneumonia among over 5	12	
Tuberculosis success rate	92%	< 90%
Were any MDR/X-TB cases diagnosed among PoCs?	Yes	No

### VACCINATION

	INDICATOR	STANDARD
Full vaccination coverage	42%	< 95%
Measles vaccination coverage		Not collected
Source of vaccination data	survey	

### IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD
Average length of stay (days)	2.8	
Case fatality rate	0.0	
Hospitalisation rate	40.8	50 - 150

## Reproductive Health

### ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	85%	> 90%
Coverage of antenatal tetanus vaccination	96%	> 95%

### DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	84%	≥ 90%
Proportion of births conducted by caesarean section	1%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	4%	< 15%
Proportion of teenage pregnancies	2%	
Still birth rate ( / 1000 total births / month)	2	
Number of maternal deaths	2	
Proportion of maternal deaths investigated within 48 hours	100%	100%

### FAMILY PLANNING

	INDICATOR	STANDARD
Contraceptive prevalence rate	2%	≥ 30%
Source of contraceptive prevalence data	HIS	

### SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD
Total number of reported rapes	0	
Proportion of eligible rape survivors provided with PEP within 72 hours		100%
Proportion of eligible rape survivors provided with ECP within 120 hours		100%

### SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD
Number of genital ulcer diseases	16	
Number of Pelvic Inflammatory diseases	482	

## HIV/AIDS

### PREVENTION & TREATMENT

Proportion of blood units screened for HIV

100%

100%

Condom distribution rate

0.01

> 0.5

Proportion of TB patients tested for HIV

100%

### PREVENTION & TREATMENT

✓ Number of PoCs receiving ART

23

✗ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

100%

100%



## Nutrition

### NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

### INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

### MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

> 70%

### CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

1383

Recovery rate

95%

> 75%

Default rate

1%

< 0%

Death rate

0%

< 10%

Referral rate

2%

### MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

2205

✓ Recovery rate

97%

> 75%

✗ Default rate

0%

< 15%

✓ Death rate

0%

< 3%

Referral rate

3%

## Water, Sanitation and Hygiene (WASH)

### WATER

Average number of litres of potable water available per person per day

< 20

Proportion of households collecting >= 15 litres per person per day

100%

> 95%

Number of persons of concern per water tap

60

< 80

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

100%

> 95%

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/100 ml

0%

> 95%

Proportion of households collecting drinking water from protected water sources only

100%

> 95%

Proportion of households with sufficient daily water storage capacity

100%

> 80%

### WATER STAFFING

Number of hygiene promoters

50

1 : < 20,000



### SANITATION

Refugees per latrine/toilet

< 20

✓ Proportion of households with drop-hole latrine or drop-hole toilet

72%

✓ Proportion of households report defecating in a toilet

94%

> 85%



✗ Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

0%

> 90%

