

Site report 2018 Ifo 2 KENYA



Population


19,894 refugees

19%

Children under 5

20%

Women of reproductive age



Origin of refugees

SOMALIA **ETHIOPIA** **DR CONGO**

Site opened
Jul 2011

Partners



Key Indicators

CRUDE MORTALITY RATE

0.0

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.1

Deaths/1,000 refugees/month

MATERNAL MORTALITY

1

Deaths

SKILLED BIRTH ATTENDANCE

86%

of all births

GAM

10%

Global Acute Malnutrition

SAM

1%

Severe Acute Malnutrition

WATER SUPPLY

35


Liters/ person/ day

LATRINE COVERAGE



4

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000
Number of qualified nurses		1 : < 10,000
Number of community health workers		1 : < 1,000

ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	55	< 50
Health utilisation rate (new visits / refugee / year)	0.5	1 - 4
Proportion of host population consultations	0%	

MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.1	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	13.1	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	13	< 20

OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%
Number of outbreaks reported	0	

MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	0	
Incidence of watery diarrhoea among children under 5	17	
Incidence of pneumonia among children under 5	22	
Incidence of pneumonia among over 5	7	
Tuberculosis success rate		< 90%
Were any MDR/X-TB cases diagnosed among PoCs?		No

VACCINATION

	INDICATOR	STANDARD
Full vaccination coverage	17%	< 95%
Measles vaccination coverage	17%	< 95%
Source of vaccination data		

IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD
Average length of stay (days)	3.1	
Case fatality rate	0.0	
Hospitalisation rate	23.6	50 - 150

Reproductive Health

ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	89%	> 90%
Coverage of antenatal tetanus vaccination	92%	> 95%

DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	86%	≥ 90%
Proportion of births conducted by caesarean section	12%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	3%	< 15%
Proportion of teenage pregnancies	5%	
Still birth rate (/ 1000 total births / month)	4	
Number of maternal deaths	1	
Proportion of maternal deaths investigated within 48 hours	100%	100%

FAMILY PLANNING

	INDICATOR	STANDARD
Contraceptive prevalence rate	6%	≥ 30%
Source of contraceptive prevalence data		

SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD
Total number of reported rapes	0	
Proportion of eligible rape survivors provided with PEP within 72 hours		100%
Proportion of eligible rape survivors provided with ECP within 120 hours		100%

SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD
Number of genital ulcer diseases	2	
Number of Pelvic Inflammatory diseases	3	

HIV/AIDS

PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%
Condom distribution rate	0.00	> 0.5
Proportion of TB patients tested for HIV		100%

PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	3		✓
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✗
PMTCT coverage	100%	100%	✓

Nutrition

NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	10%	< 20%
Prevalence of stunting (6-59 months)	20%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	1%	< 3%
Prevalence of anaemia in children (6-59 months)	52%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	46%	< 20%

INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	89%	> 80%	✗
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	88%	> 70%	✗
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	✓

MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%

CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	220	
Recovery rate	98%	> 75%
Default rate	0%	< 0%
Death rate	0%	< 10%
Referral rate	2%	

MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	228		
Recovery rate	100%	> 75%	✓
Default rate	0%	< 15%	✓
Death rate	0%	< 3%	✓
Referral rate	0%		

Water, Sanitation and Hygiene (WASH)

WATER

	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	35	< 20
Proportion of households collecting >= 15 litres per person per day		> 95%
Number of persons of concern per water tap		< 80
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%
Proportion of households collecting drinking water from protected water sources only		> 95%
Proportion of households with sufficient daily water storage capacity		> 80%

WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters		1 : < 20,000	⊖

SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	4	< 20	✓
Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of households report defecating in a toilet		> 85%	
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	