

Site report 2018 Kakuma KENYA



Population


147,966 refugees

14%

Children under 5

20%

Women of reproductive age



Origin of refugees

SOMALIA **SOUTHERN SUDAN**
ETHIOPIA

Site opened
Jan 1992

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.2 Deaths/1,000 refugees/month

MATERNAL MORTALITY

6 Deaths

SKILLED BIRTH ATTENDANCE

95% of all births

GAM

13% Global Acute Malnutrition

SAM

2% Severe Acute Malnutrition


WATER SUPPLY

21 Liters/ person/ day



LATRINE COVERAGE

7 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	4	1 : < 50,000	⚠	7	
Number of qualified nurses	36	1 : < 10,000	✓	20	
Number of community health workers	190	1 : < 1,000	✓	17	
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	78	< 50	✗	37%	< 95%
Health utilisation rate (new visits / refugee / year)	0.7	1 - 4	✗	92%	< 90%
Proportion of host population consultations	4%			Were any MDR/X-TB cases diagnosed among PoCs?	No
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	✓	Average length of stay (days)	3.3
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5	✓	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	27.8	< 30	✓	Hospitalisation rate	39.6
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	13	< 20	✓		50 - 150
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours	100%	100%	✓		
Number of outbreaks reported	1				

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	67%	> 90%	✗	72%	≥ 30%
Coverage of antenatal tetanus vaccination	86%	> 95%	⚠	Source of contraceptive prevalence data	HIS
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	95%	≥ 90%	✓	Total number of reported rapes	43
Proportion of births conducted by caesarean section	13%	5 - 15%	✓	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	8%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	54%
Proportion of teenage pregnancies	4%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	1
Number of maternal deaths	6			Number of Pelvic Inflammatory diseases	4
Proportion of maternal deaths investigated within 48 hours	100%	100%	✓		

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	✓ Number of PoCs receiving ART	469	
Condom distribution rate	0.08	> 0.5	✗ Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	99%	100% ⚠

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	13%	< 20%	⚠ Proportion of children 0-23 months who were put to the breast within one hour of birth	44%	> 80% ✓
Prevalence of stunting (6-59 months)	18%	< 20%	✓ Proportion of infants aged between 0-5 months who received only breast milk during the previous day	85%	> 70% ✓
Prevalence of severe acute malnutrition (6 - 59 months)	2%	< 3%	⚠ IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	47%	< 20%	✗ MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD
Prevalence of anaemia in women of reproductive age (15-49 yrs)	42%	< 20%	✗ Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	85%	> 70% ✓

CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	1520		Number of new moderate acute admissions (6-59 months)	3802	
Recovery rate	89%	> 75%	✓ Recovery rate	91%	> 75% ✓
Default rate	3%	< 0%	✗ Default rate	0%	< 15% ✓
Death rate	1%	< 10%	✓ Death rate	0%	< 3% ✓
Referral rate	0%		Referral rate	6%	

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	21	< 20	✓ Number of hygiene promoters	25	1 : < 20,000 ✓	
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	✓	INDICATOR	STANDARD	
Number of persons of concern per water tap	177	< 80	✗	Refugees per latrine/toilet	7 < 20 ✓	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	85%	> 95%	⚠	Proportion of households with drop-hole latrine or drop-hole toilet	35%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	✗	Proportion of households report defecating in a toilet	97%	> 85% ✓
Proportion of households collecting drinking water from protected water sources only	96%	> 95%	✓	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	88%	> 90% ⚠
Proportion of households with sufficient daily water storage capacity	89%	> 80%	✓			