

Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD	
Number of medical doctors		1 : < 50,000	•	Incidence of malaria among children under 5	0		
Number of qualified nurses		1 : < 10,000	•	Incidence of watery diarrhoea among	5		
Number of community health workers		1 : < 1,000	•	children under 5			
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	28		
Consultations per trained clinician per day	6	< 50	0	Incidence of pneumonia among over 5	3		
Health utilisation rate (new visits / refugee /	0.5	1 - 4	8	Tuberculosis success rate		< 90%	
year)				Were any MDR/X-TB cases diagnosed among	No		
Proportion of host population consultations	0%			PoCs?			
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATOR	STANDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.3	< 0.75	0	Full vaccination coverage	68%	< 95%	⊗
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5	0	Measles vaccination coverage	68%	< 95%	8
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	0	Source of vaccination data			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	0	IN-PATIENT DEPARTMENT (IPD)	INDICATOR	STANDARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate			
Number of outbreaks reported	0			Hospitalisation rate	0.0	50 - 150	8

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	100%	> 90%	v	Contraceptive prevalence rate	16%	≥ 30%	8
Coverage of antenatal tetanus vaccination	100%	> 95%		Source of contraceptive prevalence data			
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	100%	≥ 90%	0	Total number of reported rapes	6		
Proportion of births conducted by caesarean section	57%	5 - 15%	0	Proportion of eligible rape survivors provided with PEP within 72 hours	33%	100%	⊗
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	2%	< 15%	0	Proportion of eligible rape survivors provided with ECP within 120 hours	17%	100%	⊗
Proportion of teenage pregnancies	7%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	0		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receving ART	2	
Condom distribution rate	0.25	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	100%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	2%	< 20%	0	Proportion of children 0-23 months who were put to the breast within one hour of birth	30%	> 80%	⊗
Prevalence of stunting (6-59 months)	34%	< 20%	~	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	29%	> 70%	€
Prevalence of severe acute malnutrition (6 - 59 months)	0%	< 3%		IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	36%	< 20%	8	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	35%	< 20%	⊗	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%	
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)				Number of new moderate acute admissions (6-59 months)			
Recovery rate		> 75%		Recovery rate		> 75%	
Default rate		< 0%		Default rate		< 15%	
Death rate		< 10%		Death rate		< 3%	
Referral rate				Referral rate			

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	26	< 20	0	Number of hygiene promoters		1 : < 20,000	٥
Proportion of households collecting >= 15		> 95%		SANITATION	INDICATOR	STANDARD	
litres per person per day Number of persons of concern per water tap		< 80		Refugees per latrine/toilet Proportion of households with drop-hole	5	< 20	
Proportion of tests at chlorinated water		> 95%		latrine or drop-hole toilet			
collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5				Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%		Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	
Proportion of households collecting drinking water from protected water sources only		> 95%					
Proportion of households with sufficient daily water storage capacity		> 80%					