

Site report 2018 Gihembe RWANDA

Population

13,348 refugees







Partners



Key Indicators

CRUDE MORTALITY RATE

Deaths/1,000 refugees/

GAM

2% Global Acute **U5 MORTALITY RATE**

Deaths/1,000 refugees/ month

SAM

0%

Severe Acute Malnutrition

MATERNAL MORTALITY

Deaths

WATER SUPPLY

20

Liters/ person/ day

SKILLED BIRTH **ATTENDANCE**


98% of all births


LATRINE COVERAGE


Refugees/communal toilets or latrines

COMMUNICABLE DISEASES


39.0 %







NON-COMMUNICABLE DISEASES



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD		
Number of medical doctors	1	1:<50,000	Ø	Incidence of malaria among children under 5	1			
Number of qualified nurses	11	1:<10,000	②	Incidence of watery diarrhoea among	25			
Number of community health workers	30	1:<1,000	②	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	3			
Consultations per trained clinician per day	29	< 50	•	Incidence of pneumonia among over 5	2			
Health utilisation rate (new visits / refugee /	2.2	1 - 4	•	Tuberculosis success rate	100%	< 90%	②	
year) Proportion of host population consultations	0%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	©	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATOR	STAND	ARD	
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75	②	Full vaccination coverage	45%	< 95	%	8
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5	②	Measles vaccination coverage	99%	< 95	%	8
Infant Mortality Rate (IMR) (/1000 livebirths)	18.5	< 30	②	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	12	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATOR	STAND	ARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	4.1			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	54.8	50 - 1	50	②

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	30%	> 90%	8	Contraceptive prevalence rate	39%	≥ 30%	•
Coverage of antenatal tetanus vaccination	36%	> 95%	8	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	98%	≥ 90%	②	Total number of reported rapes	14		
Proportion of births conducted by caesarean section	12%	5 - 15%	②	Proportion of eligible rape survivors provided with PEP within 72 hours	50%	100%	3
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	4%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours	10%	100%	3
Proportion of teenage pregnancies	2%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	63		
Number of maternal deaths	2			Number of Pelvic Inflammatory diseases	0		
Proportion of maternal deaths investigated within 48 hours	50%	100%	8				

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	126	
Condom distribution rate	0.47	> 0.5	A	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	106%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	2%	< 20%	②	Proportion of children 0-23 months who were put to the breast within one hour of birth	92%	> 80%	•
Prevalence of stunting (6-59 months)	14%	< 20%	②	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	100%	> 70%	②
Prevalence of severe acute malnutrition (6 - 59 months)	0%	< 3%	②	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	37%	< 20%	8	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	14%	< 20%	②	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	93%	> 70%	②
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)				Number of new moderate acute admissions (6-59 months)	91		
Recovery rate		> 75%		Recovery rate	89%	> 75%	②
Default rate		< 0%		Default rate	3%	< 15%	②
Death rate		< 10%		Death rate	0%	< 3%	②
Referral rate				Referral rate	5%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	20	< 20	②	Number of hygiene promoters	25	1:<20,000	•
Proportion of households collecting >= 15 litres per person per day	50%	> 95%	8	SANITATION Refugees per latrine/toilet	INDICATOR 21	STANDARD < 20	A
Number of persons of concern per water tap		< 80		Proportion of households with drop-hole latrine or drop-hole toilet	100%		
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%		Proportion of households report defecating in a toilet	100%	> 85%	•
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%		Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%	•
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	Ø				
Proportion of households with sufficient daily water storage capacity	54%	> 80%	8				