

Site report 2018 Kigeme RWANDA



Population


20,544 refugees

14%

Children under 5

20%

Women of reproductive age



Origin of refugees

DRC

Site opened

Jun 2012

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.4 Deaths/1,000 refugees/month

MATERNAL MORTALITY

1 Deaths

SKILLED BIRTH ATTENDANCE

99% of all births

GAM

4% Global Acute Malnutrition

SAM

1% Severe Acute Malnutrition


WATER SUPPLY

13 Liters/ person/ day



LATRINE COVERAGE

38 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	1	1 : < 50,000	✓	1	
Number of qualified nurses	8	1 : < 10,000	✓	2	
Number of community health workers	22	1 : < 1,000	✓	6	
ACCESS AND UTILIZATION			MORTALITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	50	< 50	⚠	Crude Mortality Rate (CMR) (/1000/month)	0.1 < 0.75
Health utilisation rate (new visits / refugee / year)	1.5	1 - 4	✓	Under-five Mortality Rate (U5MR) (/1000/month)	0.4 < 1.5
Proportion of host population consultations	0%			Infant Mortality Rate (IMR) (/1000 livebirths)	22.4 < 30
				Neonatal Mortality Rate (NNMR) (/1000 livebirths)	14 < 20
			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
			✓	Full vaccination coverage	0% < 95% ✗
			✓	Measles vaccination coverage	99% < 95% ⚠
			✓	Source of vaccination data	HIS
			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
			✓	Average length of stay (days)	3.1
OUTBREAK ALERT AND RESPONSE				INDICATOR	STANDARD
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0
Number of outbreaks reported	0			Hospitalisation rate	33.4 50 - 150 ✗

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Antenatal care coverage	61%	> 90%	✗	43%	≥ 30% ✓	
Coverage of antenatal tetanus vaccination	97%	> 95%	✓	Source of contraceptive prevalence data	HIS	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	99%	≥ 90%	✓	Total number of reported rapes	5	
Proportion of births conducted by caesarean section	15%	5 - 15%	✓	Proportion of eligible rape survivors provided with PEP within 72 hours	100% 100% ✓	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	5%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	25% 100% ✗	
Proportion of teenage pregnancies	1%			SEXUALLY TRANSMITTED INFECTIONS (STIS)		
Still birth rate (/ 1000 total births / month)	2			Number of genital ulcer diseases	0	
Number of maternal deaths	1			Number of Pelvic Inflammatory diseases	0	
Proportion of maternal deaths investigated within 48 hours	100%	100%	✓			

HIV/AIDS

PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	1.02	> 0.5
Proportion of TB patients tested for HIV		100%

PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	187		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✓
PMTCT coverage	100%	100%	✓

Nutrition

NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	4%	< 20%
Prevalence of stunting (6-59 months)	30%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	1%	< 3%
Prevalence of anaemia in children (6-59 months)	28%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	7%	< 20%

INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	90%	> 80%	✓
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	90%	> 70%	✓
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	

MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD	
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FPF)	87%	> 70%	✓

CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)		
Recovery rate		> 75%
Default rate		< 0%
Death rate		< 10%
Referral rate		

MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	38		
Recovery rate	100%	> 75%	✓
Default rate	0%	< 15%	✓
Death rate	0%	< 3%	✓
Referral rate	0%		

Water, Sanitation and Hygiene (WASH)

WATER

	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	13	< 20	✗
Proportion of households collecting >= 15 litres per person per day	21%	> 95%	✗
Number of persons of concern per water tap		< 80	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	✓
Proportion of households with sufficient daily water storage capacity	70%	> 80%	⚠

WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters	22	1 : < 20,000	✓

SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	38	< 20	✗
Proportion of households with drop-hole latrine or drop-hole toilet	37%		
Proportion of households report defecating in a toilet	85%	> 85%	✓
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	85%	> 90%	⚠