

# Site report 2018 Kiziba RWANDA



## Population


**17,188** refugees

**11%**

Children under 5

**20%**

Women of reproductive age



Origin of refugees

**DRC**

Site opened  
Dec 1997

## Partners



## Key Indicators

CRUDE MORTALITY RATE

**0.3** Deaths/1,000 refugees/month

U5 MORTALITY RATE

**0.1** Deaths/1,000 refugees/month

MATERNAL MORTALITY

**1** Deaths

SKILLED BIRTH ATTENDANCE

**98%** of all births

GAM

**2%** Global Acute Malnutrition

SAM

**0** Severe Acute Malnutrition


WATER SUPPLY

**78** Liters/ person/ day

LATRINE COVERAGE


**6** Refugees/communal toilets or latrines

### COMMUNICABLE DISEASES




### INJURIES

### NON-COMMUNICABLE DISEASES



Total Consultations  
**42,132**

### MENTAL HEALTH



## Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	1	1 : < 50,000	✓	4	
Number of qualified nurses	7	1 : < 10,000	✓	11	
Number of community health workers	15	1 : < 1,000	⚠	25	
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	53	< 50	⚠	3%	< 95%
Health utilisation rate (new visits / refugee / year)	2.4	1 - 4	✓	99%	< 95%
Proportion of host population consultations	17%		✓	HIS	
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.3	< 0.75	✓	Average length of stay (days)	3.9
Under-five Mortality Rate (U5MR) (/1000/month)	0.1	< 1.5	✓	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	6.3	< 30	✓	Hospitalisation rate	35.9
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	4	< 20	✓		50 - 150
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%			
Number of outbreaks reported	0				

## Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	73%	> 90%	✗	59%	≥ 30%
Coverage of antenatal tetanus vaccination	91%	> 95%	⚠	HIS	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	98%	≥ 90%	✓	Total number of reported rapes	1
Proportion of births conducted by caesarean section	14%	5 - 15%	✓	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	1%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	0%
Proportion of teenage pregnancies	3%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate ( / 1000 total births / month)	0			Number of genital ulcer diseases	0
Number of maternal deaths	1			Number of Pelvic Inflammatory diseases	0
Proportion of maternal deaths investigated within 48 hours	0%	100%	✗		

## HIV/AIDS

### PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.96	> 0.5
Proportion of TB patients tested for HIV		100%

### PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	70		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✓
PMTCT coverage	96%	100%	⚠

## Nutrition

### NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	2%	< 20%
Prevalence of stunting (6-59 months)	23%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	0%	< 3%
Prevalence of anaemia in children (6-59 months)	36%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	12%	< 20%

### INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	91%	> 80%	✓
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	88%	> 70%	✓
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	

### MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD	
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/BBF)	76%	> 70%	✓

### CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)		
Recovery rate		> 75%
Default rate		< 0%
Death rate		< 10%
Referral rate		

### MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	35		
Recovery rate	88%	> 75%	✓
Default rate	7%	< 15%	✓
Death rate	0%	< 3%	✓
Referral rate	5%		

## Water, Sanitation and Hygiene (WASH)

### WATER

	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	78	< 20	✓
Proportion of households collecting >= 15 litres per person per day		> 95%	
Number of persons of concern per water tap		< 80	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5		> 95%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	✓
Proportion of households with sufficient daily water storage capacity	36%	> 80%	✗

### WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters	78	1 : < 20,000	✓

### SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	6	< 20	✓
Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of households report defecating in a toilet	100%	> 85%	✓
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%	✓