
# Site report 2018 Mahama RWANDA

### **Population**

**58,096** refugees





## **Origin of refugees**

**BURUNDI** 

Site opened Apr 2015

#### **Partners**









### **Key Indicators**

**CRUDE MORTALITY RATE** 

Deaths/1,000 refugees/

3%

**GAM** 

Global Acute

**U5 MORTALITY RATE** 

Deaths/1,000 refugees/ month

SAM

0%

Severe Acute Malnutrition

MATERNAL MORTALITY

Deaths

WATER SUPPLY

20

Liters/ person/ day

SKILLED BIRTH **ATTENDANCE** 

98% of all births

LATRINE COVERAGE


16


Refugees/communal toilets or latrines

The UN


Refugee Agency

#### COMMUNICABLE DISEASES





#### **NON-COMMUNICABLE DISEASES**



## Public Health

HEALTH STAFFING	<b>INDICATOR</b>	STANDARD		MORBIDITY	INDICATOR	STANDARD		
Number of medical doctors	5	1: < 50,000	•	Incidence of malaria among children under 5	3			
Number of qualified nurses	32	1:<10,000	<b>②</b>	Incidence of watery diarrhoea among	23			
Number of community health workers	136	1:<1,000	<b>Ø</b>	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	14			
Consultations per trained clinician per day	52	< 50	<b>A</b>	Incidence of pneumonia among over 5	5			
Health utilisation rate (new visits / refugee /	1.3	1 - 4	•	Tuberculosis success rate	100%	< 90%	<b>②</b>	
year) Proportion of host population consultations	0%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	OR STANE	DARD	
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	<b>②</b>	Full vaccination coverage	36%	< 9!	5%	8
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5	<b>②</b>	Measles vaccination coverage	100%	< 9!	5%	8
Infant Mortality Rate (IMR) (/1000 livebirths)	18.5	< 30	<b>②</b>	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	13	< 20	<b>②</b>	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STANE	DARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	4.6			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	25.4	50 -	150	8

# Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	75%	> 90%	8	Contraceptive prevalence rate	43%	≥ 30%	•
Coverage of antenatal tetanus vaccination	92%	> 95%	<b>A</b>	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	98%	≥ 90%	<b>②</b>	Total number of reported rapes	23		
Proportion of births conducted by caesarean section	4%	5 - 15%	A	Proportion of eligible rape survivors provided with PEP within 72 hours	73%	100%	3
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	4%	< 15%	<b>②</b>	Proportion of eligible rape survivors provided with ECP within 120 hours	67%	100%	3
Proportion of teenage pregnancies	0%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate ( / 1000 total births / month)	3			Number of genital ulcer diseases	77		
Number of maternal deaths	4			Number of Pelvic Inflammatory diseases	61		
Proportion of maternal deaths investigated within 48 hours	0%	100%	8				

## HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receving ART	706	
Condom distribution rate	0.39	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	99%	100%

## Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	3%	< 20%	<b>②</b>	Proportion of children 0-23 months who were put to the breast within one hour of birth	90%	> 80%	<b>②</b>
Prevalence of stunting (6-59 months)	30%	< 20%	<b>②</b>	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	83%	> 70%	<b>②</b>
Prevalence of severe acute malnutrition (6 - 59 months)	0%	< 3%	<b>②</b>	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	45%	< 20%	8	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	16%	< 20%	<b>②</b>	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	92%	> 70%	<b>②</b>
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)				Number of new moderate acute admissions (6-59 months)	551		
Recovery rate		> 75%		Recovery rate	93%	> 75%	<b>②</b>
Default rate		< 0%		Default rate	2%	< 15%	<b>②</b>
Death rate		< 10%		Death rate	0%	< 3%	<b>②</b>
Referral rate				Referral rate	3%		

# Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	20	< 20	<b>②</b>	Number of hygiene promoters	120	1:<20,000	•
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	•	SANITATION  Refugees per latrine/toilet	INDICATOR 16	STANDARD < 20	<b>o</b>
Number of persons of concern per water tap	80	< 80	•	Proportion of households with drop-hole latrine or drop-hole toilet	100%	120	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5		> 95%		Proportion of households report defecating in a toilet	100%	> 85%	•
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%		Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%	•
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	•				
Proportion of households with sufficient daily water storage capacity	95%	> 80%	•				