



Site report 2018 Abuda SUDAN



Population

4,150 refugees



Origin of refugees

ERITREA ETHIOPIA NONE

Site opened
Jan 1981

Partners



Key Indicators

CRUDE MORTALITY RATE

0.2 Deaths/1,000 refugees/month

U5 MORTALITY RATE

1.1 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

77% of all births

GAM

11% Global Acute Malnutrition

SAM

2% Severe Acute Malnutrition


WATER SUPPLY

40 Liters/ person/ day



LATRINE COVERAGE

11 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000
Number of qualified nurses		1 : < 10,000
Number of community health workers		1 : < 1,000

ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	28	< 50
Health utilisation rate (new visits / refugee / year)	3.4	1 - 4
Proportion of host population consultations	34%	

MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	1.1	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	22.2	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20

OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%
Number of outbreaks reported	0	

MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	64	
Incidence of watery diarrhoea among children under 5	173	
Incidence of pneumonia among children under 5	248	
Incidence of pneumonia among over 5	53	
Tuberculosis success rate		< 90%
Were any MDR/X-TB cases diagnosed among PoCs?		No

VACCINATION

	INDICATOR	STANDARD	
Full vaccination coverage	79%	< 95%	✗
Measles vaccination coverage	85%	< 95%	⚠
Source of vaccination data			✓

IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD	
Average length of stay (days)	1.0		
Case fatality rate	0.0		
Hospitalisation rate	263.4	50 - 150	✓

Reproductive Health

ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	99%	> 90%
Coverage of antenatal tetanus vaccination	75%	> 95%

DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	77%	≥ 90%
Proportion of births conducted by caesarean section	7%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	0%	< 15%
Proportion of teenage pregnancies	49%	
Still birth rate (/ 1000 total births / month)	2	
Number of maternal deaths	0	
Proportion of maternal deaths investigated within 48 hours		100%

FAMILY PLANNING

	INDICATOR	STANDARD	
Contraceptive prevalence rate	1%	≥ 30%	✗
Source of contraceptive prevalence data			⚠

SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD	
Total number of reported rapes	0		✗
Proportion of eligible rape survivors provided with PEP within 72 hours		100%	✓
Proportion of eligible rape survivors provided with ECP within 120 hours		100%	✓

SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD	
Number of genital ulcer diseases	0		
Number of Pelvic Inflammatory diseases	0		

HIV/AIDS

PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.00	> 0.5
Proportion of TB patients tested for HIV		100%

PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	0		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✗
PMTCT coverage	53%	100%	✗

Nutrition

NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	11%	< 20%
Prevalence of stunting (6-59 months)	45%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	2%	< 3%
Prevalence of anaemia in children (6-59 months)	35%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	28%	< 20%

INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	83%	> 80%	✗
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	81%	> 70%	✗
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	

MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/BBF)		> 70%

CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	46	
Recovery rate	80%	> 75%
Default rate	0%	< 0%
Death rate	7%	< 10%
Referral rate	9%	

MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	158		
Recovery rate	99%	> 75%	✓
Default rate	1%	< 15%	✓
Death rate	0%	< 3%	✓
Referral rate	1%		

Water, Sanitation and Hygiene (WASH)

WATER

	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	40	< 20	✓
Proportion of households collecting >= 15 litres per person per day		> 95%	
Number of persons of concern per water tap		< 80	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	
Proportion of households collecting drinking water from protected water sources only		> 95%	
Proportion of households with sufficient daily water storage capacity		> 80%	

WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters		1 : < 20,000	⊖

SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	11	< 20	✓
Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of households report defecating in a toilet		> 85%	
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	