

Site report 2018 Shagarab I II III SUDAN



Population


35,497 refugees

18%

Children under 5

20%

Women of reproductive age



Origin of refugees

ERITREA

ETHIOPIA

SOMALIA

NONE

Site opened
Jan 1985

Partners



Key Indicators

CRUDE MORTALITY RATE

0.2

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.3

Deaths/1,000 refugees/month

MATERNAL MORTALITY

0

Deaths

SKILLED BIRTH ATTENDANCE

95%

of all births

GAM

13%

Global Acute Malnutrition

SAM

2%

Severe Acute Malnutrition

WATER SUPPLY

19


Liters/ person/ day

LATRINE COVERAGE


19

Refugees/communal toilets or latrines

COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES

MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000	⊖	Incidence of malaria among children under 5	20
Number of qualified nurses		1 : < 10,000	⊖	Incidence of watery diarrhoea among children under 5	20
Number of community health workers		1 : < 1,000	⊖	Incidence of pneumonia among children under 5	224
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	62	< 50	⊗	Incidence of pneumonia among over 5	85
Health utilisation rate (new visits / refugee / year)	3.6	1 - 4	✓	Tuberculosis success rate	< 90%
Proportion of host population consultations	16%			Were any MDR/X-TB cases diagnosed among PoCs?	No
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75	✓	Average length of stay (days)	1.9
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5	✓	Case fatality rate	0.1
Infant Mortality Rate (IMR) (/1000 livebirths)	4.9	< 30	✓	Hospitalisation rate	43.9
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	✓		50 - 150
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD			
Proportion of outbreaks investigated within 48 hours		100%			
Number of outbreaks reported	0				

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	104%	> 90%	✓	Contraceptive prevalence rate	0%
Coverage of antenatal tetanus vaccination	104%	> 95%	✓	Source of contraceptive prevalence data	≥ 30%
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	95%	≥ 90%	✓	Total number of reported rapes	0
Proportion of births conducted by caesarean section	3%	5 - 15%	⊗	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	0%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%
Proportion of teenage pregnancies	26%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	0
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	8
Proportion of maternal deaths investigated within 48 hours		100%			

HIV/AIDS

PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.01	> 0.5
Proportion of TB patients tested for HIV		100%

PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	0		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✗
PMTCT coverage	11%	100%	✗

Nutrition

NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	13%	< 20%
Prevalence of stunting (6-59 months)	53%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	2%	< 3%
Prevalence of anaemia in children (6-59 months)	42%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	28%	< 20%

INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	89%	> 80%	✗
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	77%	> 70%	✗
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	

MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/BBF)		> 70%

CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	363	
Recovery rate	97%	> 75%
Default rate	0%	< 0%
Death rate	2%	< 10%
Referral rate	0%	

MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	681		
Recovery rate	99%	> 75%	✓
Default rate	0%	< 15%	✓
Death rate	1%	< 3%	✓
Referral rate	0%		

Water, Sanitation and Hygiene (WASH)

WATER

	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	19	< 20	⚠
Proportion of households collecting >= 15 litres per person per day		> 95%	
Number of persons of concern per water tap		< 80	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	
Proportion of households collecting drinking water from protected water sources only		> 95%	
Proportion of households with sufficient daily water storage capacity		> 80%	

WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters		1 : < 20,000	⊖

SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	19	< 20	✓
Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of households report defecating in a toilet		> 85%	
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	