

Site report 2018 Girba SUDAN



Population


7,486 refugees

18%

Children under 5

20%

Women of reproductive age



Origin of refugees

ERITREA

SOMALIA

NONE

Site opened
Jan 1979

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0

Deaths/1,000 refugees/month

MATERNAL MORTALITY

0

Deaths

SKILLED BIRTH ATTENDANCE

94%

of all births

GAM

9%

Global Acute Malnutrition

SAM

2%

Severe Acute Malnutrition

WATER SUPPLY

42


Liters/ person/ day

LATRINE COVERAGE


14

Refugees/communal toilets or latrines

COMMUNICABLE DISEASES




INJURIES



Total Consultations
19,176

NON-COMMUNICABLE DISEASES



MENTAL HEALTH

Public Health

HEALTH STAFFING

| | INDICATOR | STANDARD |
|------------------------------------|-----------|--------------|
| Number of medical doctors | | 1 : < 50,000 |
| Number of qualified nurses | | 1 : < 10,000 |
| Number of community health workers | | 1 : < 1,000 |

ACCESS AND UTILIZATION

| | INDICATOR | STANDARD |
|---|-----------|----------|
| Consultations per trained clinician per day | 36 | < 50 |
| Health utilisation rate (new visits / refugee / year) | 2.6 | 1 - 4 |
| Proportion of host population consultations | 24% | |

MORTALITY

| | INDICATOR | STANDARD |
|---|-----------|----------|
| Crude Mortality Rate (CMR) (/1000/month) | 0.1 | < 0.75 |
| Under-five Mortality Rate (U5MR) (/1000/month) | 0.0 | < 1.5 |
| Infant Mortality Rate (IMR) (/1000 livebirths) | 0.0 | < 30 |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths) | 0 | < 20 |

OUTBREAK ALERT AND RESPONSE

| | INDICATOR | STANDARD |
|--|-----------|----------|
| Proportion of outbreaks investigated within 48 hours | | 100% |
| Number of outbreaks reported | 0 | |

MORBIDITY

| | INDICATOR | STANDARD |
|--|-----------|----------|
| Incidence of malaria among children under 5 | 36 | |
| Incidence of watery diarrhoea among children under 5 | 28 | |
| Incidence of pneumonia among children under 5 | 103 | |
| Incidence of pneumonia among over 5 | 43 | |
| Tuberculosis success rate | | < 90% |
| Were any MDR/X-TB cases diagnosed among PoCs? | | No |

VACCINATION

| | INDICATOR | STANDARD |
|------------------------------|-----------|----------|
| Full vaccination coverage | 161% | < 95% |
| Measles vaccination coverage | 163% | < 95% |
| Source of vaccination data | | |

IN-PATIENT DEPARTMENT (IPD)

| | INDICATOR | STANDARD |
|-------------------------------|-----------|----------|
| Average length of stay (days) | 3.2 | |
| Case fatality rate | 0.0 | |
| Hospitalisation rate | 95.2 | 50 - 150 |

Reproductive Health

ANTENATAL CARE

| | INDICATOR | STANDARD |
|---|-----------|----------|
| Antenatal care coverage | 100% | > 90% |
| Coverage of antenatal tetanus vaccination | 100% | > 95% |

DELIVERY CARE

| | INDICATOR | STANDARD |
|--|-----------|----------|
| Proportion of births attended by skilled personnel | 94% | ≥ 90% |
| Proportion of births conducted by caesarean section | 12% | 5 - 15% |
| Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs) | 0% | < 15% |
| Proportion of teenage pregnancies | 0% | |
| Still birth rate (/ 1000 total births / month) | 1 | |
| Number of maternal deaths | 0 | |
| Proportion of maternal deaths investigated within 48 hours | | 100% |

FAMILY PLANNING

| | INDICATOR | STANDARD |
|---|-----------|----------|
| Contraceptive prevalence rate | 0% | ≥ 30% |
| Source of contraceptive prevalence data | | |

SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

| | INDICATOR | STANDARD |
|--|-----------|----------|
| Total number of reported rapes | 0 | |
| Proportion of eligible rape survivors provided with PEP within 72 hours | | 100% |
| Proportion of eligible rape survivors provided with ECP within 120 hours | | 100% |

SEXUALLY TRANSMITTED INFECTIONS (STIS)

| | INDICATOR | STANDARD |
|--|-----------|----------|
| Number of genital ulcer diseases | 0 | |
| Number of Pelvic Inflammatory diseases | 0 | |

HIV/AIDS

| PREVENTION & TREATMENT | INDICATOR | STANDARD | PREVENTION & TREATMENT | INDICATOR | STANDARD |
|--|-----------|----------|--|-----------|----------|
| Proportion of blood units screened for HIV | 29% | 100% | ✓ Number of PoCs receiving ART | 0 | |
| Condom distribution rate | 0.00 | > 0.5 | ✗ Proportion of HIV positive mothers who swallowed ARV during delivery | | 100% |
| Proportion of TB patients tested for HIV | | 100% | PMTCT coverage | 61% | 100% ✗ |

Nutrition

| NUTRITION GENERAL | INDICATOR | STANDARD | INFANT AND YOUNG CHILD FEEDING (IYCF) | INDICATOR | STANDARD |
|--|-----------|----------|---|-----------|----------|
| Prevalence of global acute malnutrition (6-59 months) | 9% | < 20% | ✓ Proportion of children 0-23 months who were put to the breast within one hour of birth | 93% | > 80% ✗ |
| Prevalence of stunting (6-59 months) | 30% | < 20% | ✓ Proportion of infants aged between 0-5 months who received only breast milk during the previous day | 91% | > 70% ✗ |
| Prevalence of severe acute malnutrition (6 - 59 months) | 2% | < 3% | ✓ IYCF programmes targeting pregnant and lactating women established or maintained? | | Yes |
| Prevalence of anaemia in children (6-59 months) | 38% | < 20% | ✗ MICRONUTRIENT DEFICIENCY | INDICATOR | STANDARD |
| Prevalence of anaemia in women of reproductive age (15-49 yrs) | 27% | < 20% | ✗ Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF) | | > 70% |

| CMAM: SAM TREATMENT | INDICATOR | STANDARD | MAM TREATMENT | INDICATOR | STANDARD |
|---|-----------|----------|---|-----------|----------|
| Number of new severe acute admissions (6-59 months) | 58 | | Number of new moderate acute admissions (6-59 months) | 208 | |
| Recovery rate | 98% | > 75% | ✓ Recovery rate | 95% | > 75% ✓ |
| Default rate | 0% | < 0% | ✗ Default rate | 0% | < 15% ✓ |
| Death rate | 2% | < 10% | ✓ Death rate | 0% | < 3% ✓ |
| Referral rate | 0% | | Referral rate | 4% | |

Water, Sanitation and Hygiene (WASH)

| WATER | INDICATOR | STANDARD | WATER STAFFING | INDICATOR | STANDARD |
|--|-----------|----------|---|-----------|----------------|
| Average number of litres of potable water available per person per day | 42 | < 20 | ✓ Number of hygiene promoters | | 1 : < 20,000 - |
| Proportion of households collecting >= 15 litres per person per day | | > 95% | SANITATION | INDICATOR | STANDARD |
| Number of persons of concern per water tap | | < 80 | Refugees per latrine/toilet | 14 | < 20 ✓ |
| Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5 | | > 95% | Proportion of households with drop-hole latrine or drop-hole toilet | | > 85% |
| Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml | | > 95% | Proportion of households report defecating in a toilet | | > 90% |
| Proportion of households collecting drinking water from protected water sources only | | > 95% | Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety) | | > 90% |
| Proportion of households with sufficient daily water storage capacity | | > 80% | | | |